

**FRANCE. PROGRESS REPORT ON CETACEAN RESEARCH, APRIL 2000 TO APRIL 2001, WITH
STATISTICAL DATA FOR THE CALENDAR YEAR 2001.**

Compiled by Daniel ROBINEAU

Muséum National d'Histoire Naturelle, Laboratoire d'Anatomie comparée, 55 rue Buffon, F 75005 Paris.

This report summarises information obtained from: Centre de Recherche sur les Mammifères Marins (CRMM), La Rochelle. Groupe de Recherche sur les Cétacés (GREC), Antibes. Opération Cétacés Nouvelle Calédonie (OCNC), Noumea.

1. Species and stocks studied

Common name	Scientific name	Area/stock(s)	Items referred to
Cetaceans	<i>Cetacea</i>	French continental coast	4.3, 5, 7.1, 8
Cetaceans	<i>Cetacea</i>	French continental waters	2.1.2, 3.1.1,
Cetaceans	<i>Cetacea</i>	North-East Atlantic	2.1.2,
Sperm whale	<i>Physeter macrocephalus</i>	Mediterranean Sea	2.1.1, 3.1.1, 9
Humpback whale	<i>Megaptera novaeangliae</i>	French Polynesia	2.1.1, 3.1.1
Humpback whale	<i>Megaptera novaeangliae</i>	New Caledonia	2.1.1, 2.2, 3.2, 4.1, 9
Cetaceans	<i>Cetacea</i>	New Caledonia	2.1.2, 2.2

2. Sightings data

2.1 Field work

2.1.1 SYSTEMATIC

- Sperm whale, Mediterranean Sea (GREC) : dedicated distribution survey from small boat (June-July 2000).
- Humpback whale, French Polynesia (GREC) : dedicated distribution survey from small boat (September-October 2000).
- Humpback whales, New Caledonia (OCNC). Survey held from 8 July to 10 September, 2000 : 50 observation days in the southern lagoon and 10 observation days in the northern lagoon of New Caledonia.

2.1.2 OPPORTUNISTIC, PLATFORMS OF OPPORTUNITY

- Cetaceans, French waters (CRMM). In 1999 the CRMM recorded 250 sightings of cetaceans: 200 in the Atlantic (mostly in the Bay of Biscay) and the English Channel, 50 in the Western Mediterranean Sea.
- Cetaceans, New Caledonia (OCNC). Sighting data are collected from incidental sighting forms returned by the general public (total holdings : 466 forms, analysis in progress).

2.2 Analyses/development of techniques

- Cetaceans, New Caledonia (OCNC)

Inventory of marine mammal species using opportunistic sightings collected over a 9 year period lead to a total of 13 species (Garrigue and Greaves 2001).

Estimation of the humpback whale population ranged between 243 and 386 individuals using weighted mean of Petersen estimates (Garrigue et al. 2000a). The crude birth rate is included in between 3.4 and 10 % (Garrigue et al. 2000a).

3. Marking data

3.1 Field work

3.1.1 NATURAL MARKING DATA

Species	Feature	Area/stock	Calendar year/season/no. photographed	Catalogued (Y/N)	Catalogue total	Contact person/insititute
Bottlenose d.	Dors.Fin	Atlantic	2001/6	Y	6	CRMM, La Rochelle
Long-fin. pilot w.	Dors Fin	Atlantic	2000/25	Y	31	CRMM, La Rochelle
Humpback whale	Fluke	N. Caledonia	2000/57 (27 new id.)	Y	206	OCNC
Sperm whale	Fluke	Mediterran.	2000/6	Y		GREC & NAMSWC

3.2 Analyses/ development of techniques

- Humpback whales, New Caledonia (OCNC)

Photographic comparison have been done on a regularly basis with eastern Australia (Garrigue et al. 2000b). More comparisons have been realized during two workshops of the South Pacific Whale Research Consortium. These involved catalogues of humpback whales in New Zealand, Tonga, East Australia, French Polynesia, Cook Islands, Antarctic Peninsula (INACH), part of Columbia and Ecuador. The number of resight are : East Australia : 9, New Zealand : 2, Tonga : 7, other countries or areas : 0.

4. Tissue/biological samples collected

4.1 Biopsy samples

Species	Area/stock	Calendar year/ season no. collected	Archived (Y/N)	No. analyzed	Total holdings	Contact person/institute
Humpback w.	N. Caledonia	2000/65	Y		217	OCNC & S. Baker, Univ Auckland.

4.3 Samples from stranded animals

Species	Area/stock	Calendar year/ season	Archived (Y/N)	Tissue type(s)*	Contact person/institute
Fin whale	Atlantic & Med	2000/3	Y	Various tissues	CRMM , La Rochelle
Delphinids spp	Atlantic	2000/2	Y	Various tissues	CRMM , La Rochelle
Bottlenose d.	Atlantic.	2000/7	Y	Various tissues	CRMM ; La Rochelle
Striped d.	Atlantic	2000/12	Y	Various tissues	CRMM , La Rochelle
Common d.	Atlantic	2000/126	Y	Various tissues	CRMM , La Rochelle
Long fin. pilot. w	Atlantic	2000/2	Y	Various tissues	CRMM , La Rochelle
Harbour p.	Atlantic	2000/11	Y	Various tissues	CRMM , La Rochelle
White beaked d.	Atlantic	2000/1	Y	Various tissues	CRMM , La Rochelle

*e.g. liver, skin, blubber etc.

5. Pollution studies

- Samples are collected from stranded specimens on the coasts of France by CRMM and pollutant analysis (heavy metals) are carried out by the LBME, University of La Rochelle. In 2000, 70 specimens were sampled for ecotoxicological study, including 3 Bottlenose dolphin, 7 Striped dolphins, 50 Common dolphins, 9 Harbour porpoises, 1 White beaked dolphin.

- Humpback whales, New Caledonia (OCNC). 54 blubber samples collected in 2000 (total holding: 197).

7. Statistics for small cetaceans

7.1 For the calendar year 2000

In 2000, CRMM recorded 3 incidental catches of harbour porpoises (Atlantic stock) in fishing nets and 245 stranded cetaceans with marks suggesting incidental by-catch in fishing gear.

Species	Area/stock	Directed catch		Incidental mortality			Live-capture
		Reported	Est. total	Reported	Est. total	Source	
Bottlenose dolphin	Atlantic			6		CRMM	
Striped dolphin	Atlantic			11		CRMM	
Striped dolphin	Medit.			4		CRMM	
Common dolphin	Atlantic			201		CRMM	
Common dolphin	English Ch.			7		CRMM	
Long finned pilot w	Atlantic			1		CRMM	
Long finned pilot w.	English Ch.			1		CRMM	
Harbour porpoise	Atlantic			11		CRMM	
Harbour porpoise	English Ch.			3		CRMM	

8. Strandings

In 2000, CRMM (La Rochelle), recorded 820 stranded cetaceans on the coasts of France.

9. Other studies and analyses

- Good acoustic recordings are used by GREC (Antibes), to study the stock identity of two whale species:
 - Humpback whale, French Polynesia: breeding songs (1997, 1998, 1999, 2000 data/analog and digital tapes/several individuals recorded);
 - Sperm whale, Mediterranean Sea: social vocalisations (1999, 2000 data/ analog and digital tapes/feeding and nursery schools recorded).
- Humpback whale, New Caledonia (OCNC). 7 acoustic recording of songs were collected during 1999 breeding season (total holding 57 recordings).
- Humpback whale, New Caledonia (OCNC): whale watching. Studies on commercial activity of whale watching have been carried on (Virly and Garrigue 2000, Garrigue and Virly 2000, Virly 2000)

10. Literature cited

- Garrigue, C. and J. Greaves, 2001. Cetacean records for the New Caledonian area (South West Pacific). *Micronesia*, 24 (1): 27-33.
- Garrigue, C, Greaves, J. and M. Chamberllant, 2000a. Characteristic of the New Caledonian humpback whales populations. *Humpback 2000 Conf.*, Brisbane, Australia.
- Garrigue, C; and S. Virly, 2000. Whale watching in New Caledonia: a new industry. *Humpback 2000 Conf..*, Brisbane, Australia.
- Virly, S. 2000. Activité de whale watching en Nouvelle Calédonie (saison 2000). Etude de satisfaction. *Rapport réalisé pour le Service du Tourisme de la Province sud, BP 295, 98846 Noumea, NC.* 53p.
- Virly, S. and Garrigue, C. 2000. Étude de l'activité commerciale de l'observation des mammifères marins en Nouvelle Calédonie. *Rapport réalisé pour le Service du Tourisme de la Province sud, BP 295, 98846 Noumea, NC.* : 55p.

11. Publications

11.1 Published or 'In Press' papers only

- Buffrénil, V. De, Zylberberg, L., Traub, W. and Casinos, A. 2000. Structural and mechanical characteristics of hyperdense bone of the rostrum of *Mesoplodon densirostris* (Cetacea, Ziphiidae): summary of recent observations. *Historical Biology* 14 : 57-65.
- Drouot, V., Gannier, I. and J. C. Goold, 2000. Underwater vocalisations for assessing sperm whale habitat. *14th Conf. Europ. Cetacean Soc.* (Cork, 2-5 April 2001), Abstracts.
- Duhamel, G. 2001. Extant vertebrates in marine food chains and trophic networks. In: J. M. Mazin and V. De Buffrénil (eds) *Adaptation of tetrapods to life in water*; Verlag. F. Pfeil, München:55-90.
- Fordyce, E. and Ch. De Muizon, 2001. Evolutionary history of the cetaceans: a review. In: J. M. Mazin and V. De Buffrénil (eds) *Adaptation of tetrapods to life in water*; Verlag. F. Pfeil, München: 169-234.
- Gannier, A. 2000. Distribution of cetaceans off the Society Island (French Polynesia) as obtained from dedicated

survey. *Aquatic Mammals* 26 (2): 11-126.

Garrigue, C. and J. Greaves, 2001. Cetacean records for the New Caledonian area (South West Pacific). *Micronesia*, 24 (1): 27-33.

Garrigue, C., Greaves, J. and M. Chamberllant, 2000a. Characteristic of the New Caledonian humpback whales populations. *Humpback 2000 Conf.*, Brisbane, Australia.

Garrigue, C., Forestelle, P., Greaves, J., Gill, P., Naessig, P. Baker, C. S. and N. Patenaude, 2000b. Migratory movement of humpback whales (*Megaptera novaeangliae*) between New Caledonia, East Australia and New Zealand. *J. Cetacean Res. & Manage.* 2 (2): 111-115.

Garrigue, C., Fernandez, J. M., Badie, J. M., Bernard, Ch., Greaves, J., Rivaton, J. and M. Trescinski, 2000. Impact of human activities in short-finned pilot whales (*Globicephala macrorhynchus*) and Pygmy sperm whale (*Kogia breviceps*) of the South West Pacific ocean by measuring Cs-137, K-40 and Pb210. *PHERA 2000*, June 2000, Noumea, New Caledonia.

Garrigue, C. and S. Virly, 2000. Whale watching in New Caledonia: a new industry. *Humpback 2000 Conf.*, Brisbane, Australia.

Pichler, F.B., Robineau, D., Goodall, R. N. P., Meyer, M., Olavarria, C. and C. S. Baker, 2001. Origin and radiation of Southern Hemisphere coastal dolphins (genus *Cephalorhynchus*). *Molecular Ecol.* (in Press).

Robineau, D., 2001. Distributional ecology of extant marine mammals. In: J. M. Mazin and V. De Buffrénil (eds) *Adaptation of tetrapods to life in water*; Verlag. F. Pfeil, München: 311-336.

11.2 Unpublished literature

Garrigue, C., Aguayo, A., Baker, C. S. et al. 2000a. Movements of humpback whales in Oceania, South Pacific. IWC SC/52/IA6.

Davoust, L., 2000. Analyse numérique des effectifs d'échouages de mammifères marins sur le côtes françaises : tendances, fluctuations, événements. Rapport DAA Halieutique. ENSAR, Sept 2000 : 59p.

Devez, A., Van Canneyt, O. and L. Soulier, 2000. First report on specimen of Blainville's beaked whale (*Mesoplodon densirostris*) and on specimen of Gervais' beaked whale (*Mesoplodon europaeus*) stranded on the French Atlantic coast of France. *14th Conf. of the European Cetacean Society, Cork, Ireland, April 2000. Poster.*

Poncelet, E. O., Van Canneyt, O. and J. J. Boubert, 2000. Considerable amount of plastic debris in the stomach of a Cuvier's beaked whale (*Ziphius cavirostris*) washed ashore on the French Atlantic coast of France. *14th Conf. of the European Cetacean Society, Cork, Ireland, April 2000. Poster.*

Serent, S., 2000. Élaboration de l'architecture informatique d'un futur système d'information géographique : distribution des mammifères marins du proche atlantique et paramètre environnementaux. Rapport DESS ISI, Université de La Rochelle, Sept 2000 : 50p.

Van Canneyt, O., Caurant, F., Poncelet, E. and V. Ridoux, 2000. 20 years (1980-1999) of occasional sightings compilation in the Bay of Biscay. *14th Conf. of the European Cetacean Society, Cork, Ireland, April 2000. Poster.*

Van Canneyt, O., Heintz, M. and E. Poncelet, 2000. Les échouages de mammifères marins sur le littoral français en 1999. Rapport pour le MATE, DNP, Observatoire du Patrimoine Naturel : 49p.

Virly, S. 2000. Activité de whale watching en Nouvelle Calédonie (saison 2000). Etude de satisfaction. *Rapport réalisé pour le Service du Tourisme de la Province sud, BP 295, 98846 Noumea, NC.* 53p.

Virly, S. and Garrigue, C. 2000. Étude de l'activité commerciale de l'observation des mammifères marins en Nouvelle Calédonie. *Rapport réalisé pour le Service du Tourisme de la Province sud, BP 295, 98846 Noumea, NC.* : 55p.