

Status of western gray whales off northeastern Sakhalin Island, Russia in 2012

ALEXANDER M. BURDIN¹, OLGA A. SYCHENKO¹, MAXIM M. SIDORENKO²

Contact e-mail: fewr@mac.com¹

¹Kamchatka Branch of Pacific Institute of Geography, Far East Branch - Russian Academy of Sciences Petropavlovsk-Kamchatsky, RUSSIA

²Pacific Oceanological Institute, Far East Branch, Russian Academy of Sciences, Vladivostok, RUSSIA

ABSTRACT

The western gray whale population is critically endangered and its continued ability to survive is of concern. The most recent abundance estimate (for 2012) is 155, with 95% CI = 142-165. The collaborative Russia-U.S. research program on western gray whales summering off northeastern Sakhalin Island, Russia, has been ongoing since 1995 and has produced important data that has been used to determine the conservation status of this critically endangered population. This paper reviews findings from 2012 research activities and combines such with data from previous years, in some cases ranging back to an opportunistic survey in 1994. Photo-identification research conducted off Sakhalin Island in 2012 resulted in the identification of 88 whales, including five calves and four previously unidentified non-calves. One new reproductive female was recorded in 2012, resulting in a minimum of 30 females now known to have produced a calf at some point during the study. When 2012 data are combined with results from 1994-2011, a catalog of 214 photo-identified individuals has been compiled.

KEYWORDS: WESTERN GRAY WHALE; RUSSIA; POPULATION BIOLOGY; BEHAVIOR; CONSERVATION

INTRODUCTION

Gray whales (*Eschrichtius robustus*) are recognized as comprising two populations in the North Pacific Ocean. Significant mitochondrial and nuclear genetic differences have been found between whales in the western North Pacific (WNP) and those in the eastern North Pacific (ENP) (Lang *et al.*, 2011). The ENP population ranges from calving areas in Baja California, Mexico, to feeding areas in the Bering, Beaufort, and Chukchi Seas. The WNP population feeds in the Okhotsk Sea off Sakhalin Island, Russia, and in nearshore waters of the southeastern Kamchatka Peninsula (southwestern Bering Sea).

The historical distribution of gray whales in the Okhotsk Sea greatly exceeded what is found today (Reeves *et al.*, 2008). Whales associated with the Sakhalin feeding area can be absent for all or part of a given feeding season (Bradford *et al.*, 2008), indicating they probably use other areas during the summer and fall feeding period. Some of the whales identified feeding in the coastal waters off Sakhalin, including reproductive females and calves, have also been documented off the southern and eastern coast of Kamchatka (Tyurneva *et al.*, 2010). Further, whales observed off Sakhalin have been sighted off the northern Kuril Islands in the eastern Okhotsk Sea and Bering Island in the western Bering Sea (Weller *et al.*, 2003).

Recently, mixing of whales identified in the WNP and ENP has been observed. Lang (2010) reported that two adult individuals from the WNP, sampled off Sakhalin in 1998 and 2004, matched the microsatellite genotypes, mtDNA haplotypes, and sexes (one male, one female) of two whales sampled off Santa Barbara, California in March 1995. Mate and colleagues (Mate *et al.*, 2011) satellite-tracked a whale from the WNP to the ENP in 2010/2011. Finally, photographic matches between the WNP and ENP, including resightings between Sakhalin and Vancouver Island and Laguna San Ignacio, have further confirmed use of areas in the ENP by whales identified in the WNP (Weller *et al.*, 2012; Urban *et al.* 2012). Despite this level of mixing, significant mitochondrial and nuclear genetic differences between whales in the WNP and ENP have been found (Lang *et al.*, 2011).

Although it is clear that some whales feeding off Sakhalin Island during the summer/fall migrate to the west coast of North America during the winter/spring, observations of gray whales in the WNP off Japan, Korea and China during the winter/spring suggest that not all gray whales in the WNP share a common wintering ground (Weller and

Brownell, 2012). Little is known about the current migratory routes and wintering areas in the WNP, but historical evidence indicates that the coastal waters of eastern Russia, the Korean Peninsula and Japan were part of the migratory route and that areas in the South China Sea were used as wintering grounds (Weller *et al.*, 2002; Weller and Brownell, 2012).

Western gray whales travel for thousands of miles to feed on the abundant soft-bottom benthic communities in the western part of Bering and Okhotsk Seas. This feeding period is extremely important for these slow breeding whales, and any disturbance preventing, reducing or limiting feeding can potentially lead to stress in and compromise the health of these whales. The near-shore affinity of gray whales makes them particularly vulnerable to environmental fluctuations and anthropogenic activities. For the past decade, industrial development in the coastal waters of northeastern Sakhalin, namely oil and gas development and related exploration (including seismic surveys, offshore platform installation, pipeline construction, dredging, vessel traffic) became a cause for concern, especially since the oil and gas fields overlap with the primary feeding ground of western gray whales (Blokhin and Burdin, 2001; Gailey *et al.*, 2011). Many individuals return annually to the same feeding sites off northeastern Sakhalin Island, indicating a site-specific dependence to this geographic area (Weller *et al.*, 2007). This critically important habitat is especially vital for nursing females and their calves, as female energetic requirements are increased during lactation, and calves need to be ready to separate and begin to feed on their own.

The western gray whale population is critically endangered and its continued ability to survive is of concern (Weller *et al.*, 2002a; Baillie *et al.*, 2004). Hunted to such low numbers in the mid 20th century that some thought it to be extinct, the population remains highly depleted today (Weller *et al.*, 2002a; Cooke *et al.*, 2008). The International Whaling Commission (IWC) and the International Union for Conservation of Nature (IUCN) have each expressed serious concern about the status of this population and have called for urgent measures to be taken to help ensure its protection (see Baillie *et al.*, 2004; IWC, 2004; Reeves *et al.*, 2005).

This report reviews summary findings from 2012 research activities on WNP gray whales off Sakhalin Island in the Russian Far East and integrates new information with data from previous years, in some cases ranging back to 1994. Discussion of the current status of the population and a review of threats to its continued survival, including potential impacts associated with large-scale oil and gas development activities on the summer feeding ground and entrapments in trap nets off Japan during migration, are provided herein.

METHODS

The overall consistency in research design, data collection techniques and data analysis maintained in 2012 allowed inter-annual comparisons to be made. Additional information, collected during more limited surveys off Piltun in 1994 and 1995 (Brownell *et al.*, 1997; Weller *et al.*, 1999), is also presented here to better describe inter-annual trends and facilitate a long-term interpretation for some results. Data from these 1994 and 1995 studies include gray whale photographs obtained between 7-12 September 1994 during the filming of a wildlife documentary by H. Minakuchi (for description see Weller *et al.*, 1999) and from 14-20 August 1995 during a pilot study to determine the feasibility of conducting boat- and shore-based research in the study area (Brownell *et al.*, 1997).

Study area

The study area is located near Zaliv Pil'tun (referred to as Piltun Lagoon) on the northeastern shore of Sakhalin Island, Russia (Fig. 2). The lagoon is approximately 80-90 km long and 15 km across at its widest point. A single channel connecting the inner lagoon with the Okhotsk Sea occurs at 52° 50' N and 143° 20' E, and has considerable biological influence on the surrounding marine environment. A lighthouse, near the lagoon channel, served as the base from which studies reported here were conducted. The nearshore marine environment of the study site is mostly sand substrate, characterized by a gradually sloping and broad continental shelf. Water depths within 5 km of shore are mostly less than 25 m deep. Despite the similarity of Piltun Lagoon to the coastal lagoons used during the winter by eastern gray whales off Baja California, Mexico, whales do not enter this lagoon.

Photo-identification surveys

Gray whales have distinctive body markings, such as natural coloration and pigmentation patterns, as well as scars that are unique to an individual and can be used for individual recognition. Boat-based photo-identification surveys were conducted on all good weather days during the 2012 study period. Identical methodology was employed during each survey, with the primary objective of encountering and photographically identifying as many whales as possible. Previous photo-identification data gathered in the Piltun area between 1995 and 2011 used right-side dorsal flank markings for identification (Brownell *et al.*, 1997; Weller *et al.*, 1999, 2006a), and for the sake of intra- and inter-annual reliability, we continued this methodological approach. Attempts were made to photograph the right dorsal flank of each whale, followed by efforts to photograph the left dorsal flank and fluke. The majority of whales

identified to date now have images of right and left flanks as well as ventral surface of flukes in the photo-identification catalog allowing for useful identification images to be collected from nearly any body region.

Photographic/video surveys involved slow travel in a 4.5 m outboard-powered inflatable boat. To photograph whales we used a Nikon D7000 digital camera with a 100-400 mm Nikon lens and Canon GL video camera. Measures of environmental conditions, water depth, geographic position, and group size were recorded for each group photographed.

RESULTS

Survey effort and photo-identification

Eleven photo-identification surveys, with a total of 69.7 hrs spent at sea and 48.8 hrs spent in direct observation of 78 whale groups, were conducted between 21 June and 30 August in 2012 (Table 1). In 2012 due to the poor weather conditions (heavy fogs, wind and high sea state) only four surveys were conducted in July, and five in August. Two photo-ID surveys were completed in June. Between 1994 and 2012, 214 western gray whales have been identified during 395 boat-based surveys off northeastern Sakhalin Island (Table 1). One hundred one of the whales in the photo-catalog were animals first identified as calves, while the remaining 113 whales were considered non-calves (i.e. adults or subadults). However, not all of these 214 individuals are thought to be alive (see Cooke et al., 2008).

Table 1. Annual survey effort, groups encountered, and whales identified in 1994-2012.

Year	Sampling Period	Number of Surveys	Observation Hours	Groups Encountered	Whales Identified
1994	09/07 - 09/12	1			9
1995	08/15 - 08/19	5	10.1	23	28
1997	07/09 - 09/08	22	33.4	114	47
1998	07/06 - 09/29	35	50.5	125	54
1999	06/29 - 10/13	56	122	434	69
2000	06/25 - 09/16	40	56.5	365	58
2001	06/25 - 09/25	49	101.8	448	72
2002	07/01 - 09/25	36	75.6	411	76
2003	07/15 - 09/13	22	41.7	219	75
2004	07/29 - 09/12	21	33.8	194	94
2005	07/04 - 09/09	20	40.9	160	93
2006	07/23 - 08/25	10	24.1	96	79
2007	07/26 - 09/09	20	32.2	187	83
2008	07/08 - 08/21	12	47.0	38	45
2009	06/24 - 08/26	17	67.0	126	82
2010	08/09 - 08/26	4	11.5	40	42
2011	06/28 - 08/26	14	32.7	83	82
2012	06/24 - 08/30	11	48.8	78	88
Overall		395	829.6	3141	214 ¹

¹ The number of whales identified annually includes resightings of individuals from previous years, resulting in a total of 214 identified individuals. The number of whales identified does not correspond to the size of the population.

Eighty-eight naturally marked individual whales, including 5 calves, were identified during 2012 (Table 1, 2). This is less than in 2011, when the highest number of calves (12) were identified off northeastern Sakhalin among all

years of our research. By the end of August, 3 mother -calves pairs were separated, while 2 of them were still together (Table 3). Of the 83 non-calves identified in 2012, 79 whales (95.2%) had previous sightings in the Piltun area during 1994-2011 photographic efforts (Table 2). The mean pod size for all groups (n=78) encountered during 2012 was 2.3 ± 1.32 ranging from 1 to 13 individuals per pod. Feeding whales were distributed in water depths ranging from 3.7 (mostly mother-calf pairs) to 21.7 m (average 7.2 ± 2.53).

Fifty-two (59%) individual gray whales were observed only once during the 2012 field season; 22 (25%) whales were sighted twice. One whale was seen 6 times within the 2012 season. This whale, named "Bud" was female #11 in our catalog. She is a well-known whale and long-time visitor to Piltun. She has been seen most years from 1997-2012 with the exception of 2003, 2009, and 2010. She is often sighted several times per season. We first observed her to have a calf in 2002, then again in 2006, 2008, and 2011. Her 2006 calf is the young whale that was fatally entrapped off Japan in January 2007 (see Weller et al., 2008). Although she was satellite tagged in 2011, her tag stopped transmitting while she was still off Piltun during the summer feeding period.

Table 2. Annual sighting trends and resighting percentages, 1994-2012.

Year	Whales Identified	Number of Calves	New Non-Calves	% Non-Calves Previously Identified
1994 ¹	9			
1995 ¹	28	2	20	23.1%
1997	47	2	25	44.4%
1998	54	8	5	89.1%
1999	69	3	12	81.8%
2000	58	3	3	94.5%
2001	72	6	6	90.9%
2002	76	9	3	95.5%
2003	75	11	2	96.9%
2004	94	8	3	96.5%
2005	93	6	4	95.4%
2006	79	4	3	96.0%
2007	83	9	2	97.3%
2008	45	3	0	100.0%
2009	82	7	2	97.6%
2010	42	3	1	97.4%
2011	82	12	1	98.6%
2012	88	5	4	95.2%

¹ Data from 1994 and 1995 were opportunistic and pilot in nature (respectively) and are thereby viewed as incomplete for some of the reported values.

Mother-calf pairs

Five mother-calf pairs were identified in 2012. All five mothers have been sighted in the study area prior to 2012; however, one of them has never been observed in previous years with a calf. This new mother contributes to the total of 30 known reproductive females documented since 1995.

The first sighting of a mother-calf pair in 2012 occurred during our second photo-ID survey, on 29 June. Three mother-calf pairs were identified in July, and the last mother-calf pair was initially observed in August. Of all five pairs, three separated by the end of August. In two other pairs, calves remained with their mothers, and therefore, no information on weaning time for them was documented (Table 3).

Table 3. Mother -calf pair IDs, dates of first sightings and separation dates

Mother ID #	Calf ID #	Calf Nickname	First Time Observed	Weaning Date
15	202	Smart	29 June 12	NA
40	203	Silver	18 July 12	20 August 12
22	204	Milkyway	18 July 12	20 August 12
81	205	Mi-mi-mi	18 July 12	NA
38	207		20 August 12	25 August 12

DISCUSSION

A number of biological parameters in concert with a variety of human-related threats, as identified during the current long-term study and discussed below, raise concern about the ability of the western gray whale population to rebound from its highly depleted state and highlight the importance of continuing the long-term research and monitoring program.

Population size

Given to the new data about movements of some WNP gray whales from the Sakhalin feeding grounds to Mexican wintering grounds, a reevaluation of the western gray whale population estimate is necessary. Data we collected over more than 15 years research supports the idea, that the size of the western gray whale population is extremely small compared to most other baleen whale populations. Photo-identification studies off northeastern Sakhalin Island have identified a total of 214 individual whales during 395 surveys conducted between 1994 and 2012. Although our photo-catalog now contains 214 whales, not all of these individuals are assumed to be alive. A population assessment by Cooke et al. using a Bayesian individually based stage-structured model fitted to the same photo-identification data as used in the mark-recapture studies, has recently been completed. The most recent WNP gray whale abundance estimate (for 2012) is 155, with 95% CI = 142-165 (IUCN, 2012).

Reproduction and survival

In 2012 a new reproductive female was observed in the Piltun area, making the total number of known reproductive females became 30. But compared to other species and populations of large cetacean, the number of breeding females in the WNP gray whale population is still very low and the Piltun area still remains the main area of feeding for females and calves. Although calves are being born annually, the limited number of known reproductive females in combination with relatively low calf survival is likely to be limiting potential population growth (Bradford et al., 2006; IISG, 2006; Cooke et al., 2008). In recent years, the calving interval in the western population appears to be shifting from a three-year interval to a two-year interval (Weller et al., 2009). If this change persists, the general increase in calf production will continue and, in turn, contribute to an increase (albeit slow) in the growth rate of the population.

Mother-calf pairs

Five mother-calf pairs were identified during the 2012 season. All five mothers have been sighted in the study area prior to 2012; however, one of them has never been observed in previous years with a calf. The annual return of reproductive females while pregnant, resting and lactating indicates that the near shore Sakhalin Island feeding area is of significant importance to the continued survival of this population. The behavior of these females indicates that this feeding ground is vital to population survival and growth.

Threats to the population

Considering the recently documented migration of some part of whale identified off Sakhalin to the Mexican wintering areas, new threats for the WNP population appear. First of all, there is a risk that the WNP population size is even smaller than previously estimated (see Weller and Brownell 2012). Further, the migration distance from the WNP to the ENP has increased by almost 3000-4000 km. These changes create new risks for calves traveling with mothers to feeding grounds in the WNP. In addition to the biological difficulties (e.g., small population size, low number of reproductive females) that western gray whales face, the onset of large-scale oil and gas development programs off Sakhalin Island in the mid-1990s introduced new threats to the future survival of the population (Weller et al., 2002a; Reeves et al., 2005; IISG, 2006). Sakhalin Island is a region rich with large reserves of offshore oil and gas that, until recently, have been unexploited. Industrial activities on the continental shelf of this region have steadily increased in the past ten years and are scheduled to expand at a rapid pace into the future. Oil and gas development activities that may negatively impact western gray whales include: (1) disturbance from underwater noise associated with seismic surveying (Weller et al., 2002b; 2006b, 2006c), pipeline dredging, ship and helicopter traffic and plat-

form operations; (2) direct interactions between whales and an oil spill or other waterborne chemicals, ships, and possible entanglements in cables or lines; and (3) habitat changes related to seafloor modifications associated with dredging and sand pumping activities that may adversely impact gray whale prey (for reviews see Reeves et al., 2005; IISG, 2006).

It is also plausible, however, that the change reflected whales being displaced from the feeding area or, worse, indicates partial abandonment of what has traditionally been a critical feeding habitat (especially for mother-calf pairs) for the population. While natural variation in food resources and other biological factors are being investigated by industry-sponsored research groups, additional investigations need to be undertaken to examine the possible contributions of pile driving activities and a seismic survey that both occurred in close proximity to the nearshore feeding ground in summer 2008. Until more conclusive explanations can be drawn with regard to the low number of whales observed in 2008, the influence of industrial activities cannot be ruled out as contributing factors.

Another significant threat to the western gray whale population involves incidental catches in coastal net fisheries, particularly off Japan, within their migratory route (Weller et al., 2002a; Kato et al., 2005, 2006, 2007; Brownell et al., 2007; Weller et al., 2008). In 2005, three female western gray whales (one mother-calf pair and one yearling) died in fishing nets on the Pacific coast of Japan during their northward migration. Unfortunately, in 2007 another young female western gray whale died after being entrapped in a trap net also on the Pacific coast of Japan (Anonymous, 2007a,b,c; Brownell et al., 2007; Kato et al., 2007; Weller et al., 2008). Projections from recent population assessments suggest that if this level of net-related mortality continues, there is a high probability the population will decline to extinction (Cooke et al., 2008). In addition, an analysis of anthropogenic scarring of western gray whales found that 18.7% ($n = 28$) of 150 individuals identified between 1994 and 2005 were determined to have been previously entangled in fishing gear (Bradford et al., 2009), further highlighting the overall risks coastal fisheries pose to western gray whales. Finally, while nothing is known about net entrapments or entanglements in other regions (e.g., Korea and China) within the range of the population, it is likely that coastal net fisheries outside of Japan also contribute to some level of mortality.

Although there are historical data on gray whales sightings in waters off Japan, South Korea and China, and also documented net entrapments near Japan, the wintering grounds for western population remain unclear (Weller et al. 2008). Some western gray whales are seen near Kamchatka, and both intra-annual and intra-seasonal exchange of individuals between feeding grounds off Sakhalin and Kamchatka are documented (Tyurneva et al. 2010, our data). In 2010 a satellite-tagging project was initiated off northeastern Sakhalin and continued in 2011. A male gray whale, which was observed as a calf in 1997 and sighted in most years of study in the Piltun area, was tagged and tracked during October 2010–February 2011. Five more whales were tagged in 2011, but only two of them transmitted for long enough to reveal anything about migration. Both whales moved at the same time to the east the Okhotsk and Bering Sea towards North America. One of the tags (whale “Agent”) stop transmitting in the Gulf of Alaska, while the tag attached to female “Varvara” transmitted all way to Mexico lagoons and back to Sakhalin Island (Piltun area). This individual traveled over 22,000 km from Sakhalin feeding grounds to the Mexico and back in 5 months. Thirteen more whales observed on the feeding grounds off Sakhalin in different years were confirmed with photographic matches to the whales from eastern North Pacific catalog (Weller et al. 2012, Urban et al., 2012). This highlights that the range and potential threats this population may face may be on a larger scale than previously anticipated.

Such a wide range in distribution makes whales vulnerable to other unknown threats to the western gray whale population including continued mortality from an undetermined level of suspected poaching in the central portion of the range (Brownell and Kasuya, 1999; Baker et al., 2002), as well as a potential increase in the likelihood of disturbance, exposure to pollution, and probability of ship strikes due to substantial nearshore industrialization and shipping congestion throughout the migratory corridor(s).

CONCLUSIONS

Based on the results reported here, it is clear that the western gray whale population could be even smaller than we have estimated and it is precariously balanced between survival and extinction. In addition to the variety of biological factors that may be limiting population growth, large-scale oil and gas development programs that may alter the prey base or introduce disturbance to feeding whales, as well as entrapment and entanglement in fishing gear, especially in trap nets off Japan, are of serious concern with regard to the future survival of the population.

Given the continued uncertainty regarding the ability of the western gray whale population to increase from its depleted state, impacts from oil and gas development activities off the northeastern Sakhalin Island coast need to be closely monitored and stringently mitigated to reduce disturbance to the lowest possible level. In addition, net entrapments of western gray whales off Japan and possibly elsewhere can lead the population to extinction (IISG,

2006; Cooke et al., 2008; Brownell et al., 2007; Weller et al., 2008). Thus, human related mortality during migration and in the (yet to be determined) wintering area(s) must be addressed and mitigated to the lowest possible level. Where scientific knowledge is lacking, the precautionary principle should be applied as the best measure of protection. With this in mind, the photo-identification and genetic biopsy research conducted since 1995, and reviewed here, must be continued to further monitor survival of individuals, describe the overall population trend and to recommend further conservation and protection measures.

In conclusion, protection of the Sakhalin Island feeding habitat, including the coastal lagoon systems that appear integrally related to the high benthic biomass used by the whales in the nearshore area, is clearly paramount to successful conservation of the western gray whale population. The unique method of benthic feeding by these whales makes them an "umbrella" species (Hooker and Gerber, 2004), whereby protection of their habitat provides protection for the biological diversity of the entire northeastern Sakhalin Island shelf. Thus, the feeding habitat of the western gray whale needs to be considered a "hot spot" for conservation planning now and in the future and every effort should be taken to protect its biological integrity. In continuation of this research and looking for the development of additional western gray whale conservation measures, the next step should be intensifying research of gray whales off Kamchatka, and create the observation point on gray whale migration along Kamchatka coast, possibly on the lighthouse in Petropavlovsk-Kamchatsky.

ACKNOWLEDGEMENTS

A number of new young researchers joined the western gray whale project in 2012. We were fortunate to work with a wonderful cast of characters during the 2012 field season, with special thanks to Oksana Savenko, Alexey Shipulin, Stepan Varlamov, Igor Usoltsev, Alexandr Likhonov. The International Fund for Animal Welfare (IFAW) and Ocean Park Conservation Foundation Hong Kong (OPCFHK) generously provide research grants in support of this research.

LITERATURE CITED

- Anonymous. 2007a. Japanese nets threaten grey whale's survival. [News in Brief] *Nature* 445:577.
- Baillie, J.E.M., Hilton-Taylor, C. and Stuart, S.N. (Editors) 2004. 2004 IUCN Red List of Threatened Species. A Global Species Assessment. IUCN, Gland, Switzerland and Cambridge, UK. xxiv + 191 pp.
- Baker, C.S., Dalebout, M.L. and Lento, G.M. 2002. Gray whale products sold in commercial markets along the Pacific coast of Japan. *Mar. Mamm. Sci.* 18:295-300.
- Blokhin, S. A. and Burdin, A. M. 2001. Distribution, abundance and some traits of behavior of the gray whale *Eschrichtius robustus* of the Korean stock at the northeastern coast of Sakhalin. *Biologiya Morya* 27(1):15-20.
- Bradford, A.L., Weller, D.W., Wade, P.R., Burdin, A.M., Brownell, R.L., Jr. 2008. Population abundance and growth rate of western gray whales *Eschrichtius robustus*. *Endangered Spe. Res.* 6(1):1-14
- Brownell, R.L., Jr., Blokhin, S.A., Burdin, A.M., Berzin, A.A., LeDuc, R.G., Pitman, R.L. and Minakuchi, H. 1997. Observations on Okhotsk-Korean gray whales on their feeding grounds off Sakhalin Island. *Rep. Int. Whal. Commn.* 47:161-162.
- Brownell, R.L., Jr. and Kasuya T. 1999. Western gray whale captured off western Hokkaido, Japan. Paper SC/51/AS25 presented to the IWC Scientific Committee. 7 pp.
- Brownell, R.L., Jr., Kasuya, T. and Weller, D.W. 2007. Entrapment of western gray whales in Japanese fishing gear: Population threats. Paper SC/59/BRG38 presented to the IWC Scientific Committee. 9pp.
- Cooke, J., Weller, D.W., Bradford, A.L., Burdin, A.M. and Brownell, R.L, Jr. 2008. Population assessment of western gray whales in 2008. Paper SC/60/BRG11 presented to the IWC Scientific Committee. 10pp.
- Cooke, J.G. 2010. Joint population assessment of western gray whales using data from IBM and Russia-US photo-identification teams collected off Sakhalin Island through 2008. In: Report to the Western Gray Whale Advisory Panel. 23 pp.
- Gailey, G., Sychenko, O. and Würsig, B. 2011. Patterns of western gray whale behavior, movement and occurrence off Sakhalin Island, 2010. In: Report to the Western Gray Whale Advisory Panel. Western gray whale research and monitoring program in 2010, Sakhalin Island, Russia. Vol.II:4/1-4/63.
- Interim Independent Scientists Group (IISG) 2006. Report of the Interim Independent Scientists Group (IISG) on mitigation measures to protect western gray whales during Sakhalin II construction operations in 2006, Vancouver, British Columbia, 3-5

April 2006. International Union for Conservation of Nature (IUCN), Business and Biodiversity Program, [Available from <http://www.iucn.org>].

International Whaling Commission. 2004. Resolution on western north Pacific gray whale. Ann. Rep. Int. Whaling Comm. 2004.

IUCN 2012. Report of the 11th Meeting of the Western Gray Whale Advisory Panel. Geneva, Switzerland. [Available from http://www.iucn.org/wgwap/publications_and_reports/]

Kato, H., Ishikawa, H., Mogoe, T. and Bando, T. 2005. Occurrence of a gray whale, *Eschrichtius robustus*, in Tokyo Bay, April-May 2005, with its biological information. Paper SC/57/BRG18 presented to the IWC Scientific Committee.

Kato, H., Ishikawa, H., Bando, T., Mogoe, T. and Moronuki, H. 2006. Status of conservation and researches on the western gray whale in Japan, June 2005-April 2006. Paper SC/58/O14 presented to the IWC Scientific Committee.

Kato, H., Ishikawa, H., Goto, M., Miyashita, T. and Moronuki, H. 2007. Status report of conservation and researches on the western gray whale in Japan, June 2006-April 2007. Paper SC/59/O18 presented to the IWC Scientific Committee.

Laake, J., Punt, A., Hobbs, R., Ferguson, M., Rugh, D. and Breiwick, J. 2009. Re-analysis of gray whale southbound migration surveys 1967-2006. NOAA Technical Memorandum. NMFS-AFSC-203. 55pp.

Lang A.R., Weller D.W., LeDuc R., **Burdin A. M.**, Pease V.L., Litovka D., Burkanov V., Brownell R.L., Jr. 2011. Genetic analysis of stock structure and movements of gray whales in the eastern and western North Pacific. presented to the IWC Scientific Committee. SC/63/BRG10

Lang A.R., Weller D.R., LeDuc R.G., **Burdin A.M.** Brownell R.L. Jr. 2010. Genetic differentiation between western and eastern gray whale populations using microsatellite markers. presented to the IWC Scientific Committee. SC/62/BRG www.iwcoffice.org/documents/sci_com/SC62docs/SC-62-BRG8.pdf

Lang, A.R., Weller, D.W., Leduc, R.G., Burdin, A.M. and Brownell, R.L., Jr. 2005 Genetic assessment of the western gray whale population: current research and future directions. SC/57/BRG14 presented to the IWC Scientific Committee.

LeDuc, R.G., Weller, D.W., Hyde J., Burdin, A.M., Rosel, P. E. et al. 2002. Genetic differences between western and eastern gray whales (*Eschrichtius robustus*). *J. Cetacean Res. Manage.* 4(1):1-5.

Reeves, R.R., Brownell, R.L., Burdin, A., Cooke, J.C., Darling, J.D., Donovan, G.P., Gulland, F.M.D., Moore, S.E., Nowacek, D.P., Ragen, T.J., Steiner, R.G., VanBlaricom, G.R., Vedenev, A. and Yablokov, A.V. 2005. Report of the Independent Scientific Review Panel on the Impacts of Sakhalin II Phase 2 on Western North Pacific Gray Whales and Related Biodiversity. IUCN, Gland, Switzerland and Cambridge, UK. 123pp [Available from <http://www.iucn.org>]

Tyurneva O. Yu., Yakovlev Yu. M., Vertyankin V. V., Selin N. I. 2010. The peculiarities of foraging migrations of the Korean-Okhotsk gray whale (*Eschrichtius robustus*) population in russian waters of the Far Eastern seas. *Rus. Jour. of Marine Biol.* Vol. 36. No 2: 117-124

Urbán J. R., D. Weller, O. Tyurneva, S. Swartz, A. Bradford, Y. Yakovlev, O. Sychenko, H. Rosales N., S. Martínez A., A. Burdin and A. Gómez-Gallardo U. 2012. Report on the photographic comparison of the western and mexican gray whale. Paper SC/64/BRG13 presented to the IWC Scientific Committee.

Weller, D.W., Würsig, B., Bradford, A.L., Burdin, A.M., Blokhin, S.A., Minakuchi, H. and Brownell, R.L., Jr. 1999. Gray whales (*Eschrichtius robustus*) off Sakhalin Island, Russia: seasonal and annual patterns of occurrence. *Mar. Mamm. Sci.* 15:1208-1227.

Weller, D.W., **Burdin**, A.M., Ivashchenko, Y.V., Tsidulko, G.A., Brownell, R.L., Jr. 2003. Summer sightings of western gray whales in the Okhotsk and western Bering Seas. Paper SC/55/BRG9 submitted to the International Whaling Commission.

Weller, D.W., Burdin, A.M., Würsig, B., Taylor, B.L. and Brownell, R.L., Jr. 2002a. The western Pacific gray whale: a review of past exploitation, current status and potential threats. *J. Cetacean Res. Manage.* 4(1):7-12.

Weller, D.W., Ivashchenko, Y.V., Tsidulko, G.A., Burdin, A.M. and Brownell, R.L., Jr. 2002b. Influence of seismic surveys on western gray whales off Sakhalin Island, Russia in 2001. Paper SC/54/BRG14 presented to the IWC Scientific Committee. 15 pp.

Weller, D.W., Bradford, A.L., Tsidulko, G.A., Ivashchenko, Y.V., Lang, A.R., Kim, H.W., Burdin, A. M. and Brownell, R.L., Jr. 2006a. A catalog of photo-identified western gray whales from Sakhalin Island, Russia. Paper SC/58/BRG2 presented to the IWC Scientific Committee [CD available on request].

Weller, D.W., Rickards, S.H., Bradford, A.L., Burdin, A.M. and Brownell, R.L., Jr. 2006b. Influence of 1997 seismic surveys on the behavior of western gray whales off Sakhalin Island, Russia. Paper SC/58/E4 presented to the IWC Scientific Committee. 12 pp.

Weller, D.W., Tsidulko, G.A., Ivashchenko, Y.V., Burdin, A.M. and Brownell, R.L., Jr. 2006c. A re-evaluation of the influence of 2001 seismic surveys on western gray whales off Sakhalin Island, Russia. Paper SC/58/E5 presented to the IWC Scientific Committee. 8 pp.

Weller, D.W., Bradford, A.L., Lang, A.R., Kim, H.W., Sidorenko, M. et al. 2007. Western gray whales off Sakhalin Island, Russia: a joint Russia-U.S. scientific investigation July-September 2007. Interim report to the Western Gray Whale Advisory Panel. 15 pp.

Weller, D.W., Bradford, A.L., Kato, H., Bando, T., Ohtani, S., Burdin, A.M. and Brownell, R.L., Jr. 2008. Photographic match of a western gray whale between Sakhalin Island, Russia, and Honshu, Japan: First link between feeding ground and migratory corridor. *J. Cetacean Res. Manage.* 10(1):89-91.

Weller, D.W., Bradford, A.L., Lang, A.R., Burdin, A.M. and Brownell, R.L., Jr. 2009. Birth intervals and sex composition of western gray whales summering off Sakhalin Island, Russia. Paper SC/61/BRG10 presented to the IWC Scientific Committee. 7 pp.

Weller, D.W., Klimek, A., Bradford, A.L., Calambokidis, J., Lang, A.R., Gisborne, B., Burdin, A.M., Szaniszló, W. and Brownell, R.L., Jr. 2011. Movements of western gray whales from the Okhotsk Sea to the eastern North Pacific. Paper SC/63/BRG6 presented to the IWC Scientific Committee. 5 pp.

Weller, D.W., Klimek, A., Bradford, A.L., Calambokidis, J., Lang, A.R., Gisborne, B., Burdin, A.M., Szaniszló, W., Urbán, J., Gomez-Gallardo Unzueta, A., Swartz, S. and Brownell, R.L., Jr. 2012. Movements of gray whales between the western and eastern North Pacific. *Endangered Species Research*, 18:193-199 doi: 10.3354/esr00447

Weller, D.W. and Brownell, R.L., Jr. 2012. A re-evaluation of gray whale records in the western North Pacific. Paper SC/64/BRG10 presented to the International Whaling Commission Scientific Committee [Available from <http://www.iwcoffice.org/>]

Zhu, Q. 2012. Gray whale bycaught in Pingtan, China. *Cetoken Newsletter* No. 29, 2012.2.1