# SC/67B/CMP/02

### Density-dependent changes in the distribution of southern right whales (Eubalaena australis) in the breeding ground Peninsula Valdes.

Nicolas Sueyro, Enrique Crespo, Magdalena Arias, Mariano Coscarella



Papers submitted to the IWC are produced to advance discussions within that meeting; they may be preliminary or exploratory. It is important that if you wish to cite this paper outside the context of an IWC meeting, you notify the author at least six weeks before it is cited to ensure that it has not been superseded or found to contain errors.

## Density-dependent changes in the distribution of southern right whales (*Eubalaena australis*) in the breeding ground Peninsula Valdés.

#### Nicolás Sueyro<sup>1,2</sup>; Enrique Crespo<sup>1,2</sup>; Magdalena Arias<sup>3</sup>; Mariano Coscarella<sup>1,2\*</sup>

1- Centro Para de Estudios de Sistemas Marinos. Blvd. Alte. Brown 2915, Puerto Madryn (Chubut), Argentina. Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET)

2 Universidad Nacional de la Patagonia San Juan Bosco - Fac. Cs. Nat. Sede Puerto Madryn, Blvd. Brown 3051

3 - CIMAS, Escuela Superior de Ciencias Marinas - Universidad Nacional del Comahue. San Martín 247 (8520) San Antonio Oeste (Río Negro), Argentina

\*Corresponding author: coscarella@cenpat.edu.ar

#### Abstract

Southern Right Whales (*Eubalaena australis*) is experiencing a population growth in the South-western Atlantic Ocean. In the breeding ground of Peninsula Valdés, as a consequence of the population increase, expansion to new areas and a change in the habitat use of the coastal area were recorded. Also, as a consequence of population increase changes in distribution and density in the core area are suspected. We analyzed information that was gathered from aerial surveys developed along the coast of Peninsula Valdés for 19 years. These surveys were divided into 4 periods. A geographical analysis of 620 km of coast revealed that in 5 km-length segments the density of whales increased to a maximum near to 3 whales per km<sup>2</sup>. This figure is proposed as a threshold that elicits a density dependence response, where the *Mother-calf* pairs remain in the area, while the *other* groups decreased their density, forcing them to move to other areas.

#### Introduction

The Southern Right Whale (*Eubalaena australis*) has a circumpolar distribution in the southern hemisphere. This species was the object of a commercial exploitation between the XVIII and XX centuries that put the species on the brink of extinction (Richards 2009). The species was protected for the first time in 1936 and additionally, in 1986, the moratorium on commercial catch established by the International Whaling Commission (IWC) came into force. By the mid-1970's several populations have shown evidence of recovery, with a doubling time of 10 to 12 years (Bannister 2001, Best et al. 2001, Cooke

et al. 2001). However, there are other populations that are still very small and there is uncertainty about their recovery (Galletti Vernazzani et al. 2014). Even today, the geographical distribution of all breeding and feeding areas of the species prior to its exploitation is unknown (IWC 2001).

The estimated population size for the species in 1997 was 7,500 animals (including 547 mature females, from Argentina and 659 from South Africa), with an estimated average growth rate for all populations in the Southern Hemisphere of 7.5% (IWC 2012). The breeding population of Peninsula Valdés is one of the best studied, with a long-term research program carried out in the Peninsula Valdés area since 1970 (Payne 1986). The population size and other parameters derived from capture-recapture models have been estimated, based on the individual recognition of the whales. The population growth rate was estimated to be around 8% at that time (Payne et al. 1981, Payne 1986, Whitehead et al. 1986, Payne et al. 1990), and recent studies estimate a 5.1% for 2010 (Cooke 2012) and 6.5% for 2012 (Cooke et al. 2015) using the same techniques.

Each year, between August and September, the number of whales reaches its maximum in the area (Crespo et al. 2014). In 1999, during the period of maximum concentration, 556 whales of all ages and classes were counted, whereas for 2007 this number increased to 1006, and in 2013, 1556 animals were reached. In 1999, 176 calves were counted, compared to 317 in 2007 and 466 in 2013 (Crespo et al. 2015). The authors estimated for the year 2007 an increasing rate of 6.22%. For the year 2014, the models indicated that the population grew at a rate of 3.23% per year. This is an indication of a decrease at a rate of -0.45% per year in the rate of population increase (Crespo et al. 2015). However, the reduction in the rate of increase was not uniform across the different groups that comprise the population. The estimate of the growth rate for the number of offspring born in the Peninsula Valdés was 5.54% per year between 1999 and 2014 (Crespo et al. 2015), almost doubling the rate of increase for the whole population during the same period. However, the rate of increase of the other class groups (namely Solitary Individuals and Breeding groups) is now close to 0% (Crespo et al. 2015). Mother-calf pairs have displaced the Other groups from the shore, and the Solitary Individuals and the Breeding groups are now, or in deeper waters far away from the coast, or have moved away from the area. This change in Peninsula Valdés distribution could mean that the coastal zone (within 2km from shore) is close to its carrying capacity (Crespo et al. 2015). Therefore, in spite of the overall decrease in the rate of increase observed in the sampling area, it is proposed that the population of the south-western Atlantic is still growing at a rate that

would be the combination of its growth in the Peninsula Valdés area and the rate occupation of new zones (Crespo et al. 2015). If this scenario is correct, whales should begin to move to other less dense regions (*ie:*, sub-optimal habitats) where the growth rate should be higher (Verner et al. 1986, Hobbs and Hanley 1990). There are different indicators support this hypothesis for Peninsula Valdés as evidenced by the number of whales occupying areas with deeper waters in Peninsula Valdés and the growing number of whales observed in Golfo San Matías, Buenos Aires, Uruguay and Santa Catarina in the southern Brazil (Groch et al. 2005, IWC 2011, Crespo et al. 2014).

The expansion of whales to other areas may be determined by the priority occupation of mothers with offspring in optimal habitat areas (Barendse and Best 2014, Carroll et al. 2014, Danilewicz et al. 2016). If this is the case, there should be a threshold density in which the *Other* groups increase their density in adjacent areas (*eg:* nonpreferred by *Mother-calf* pairs).

#### Methods

The censuses were carried out from a single-engine high-wing CESSNA B-182 aircraft, flying at a constant height of 500 feet (152 m) and at 80/90 knots every 45 days from April to December each year (Crespo et al. 2011). In each survey a distance of 620 km was covered in 5 hours of flight, flying from south to north along the coast. The surveyed area (Figure 1) is located between the mouth of the Rio Chubut and Puerto Lobos on the border with the province of Rio Negro. The width of the strip is composed by 500 meters from the coast plus approximately 1000 meters from the plane to the open sea, composing a surveyed strip of 1500 meters (Crespo et al. 2011). This strip is set to cover the "whale-road" as described by Payne (1986), where more than 90% of the whales in the area concentrate near the coast in shallow waters.



Figure 1: Sampling area. The thick black line along the coast represent the surveyed area.



Figure 2: Schematic of the survey.

The team comprised a pilot, a recorder sitting next to the pilot and two observers in the rear seats, one on the left and one on the right side of the aircraft (Crespo et al. 2014). The observations were made with the naked eye, and the information was recorded in

spreadsheets or tablet applications developed *ad-hoc* with Cyber Tracker <sup>TM</sup>. Information on the on the group composition was recorded, including *Mother-calf* pairs, *Solitary Individuals* or *Breeding* groups comprising n-1 males and one female (Crespo et al. 2011). *Solitary Individuals* and *Breeding* groups were pooled in *Other* groups, as a group category opposite to the *Mother-calf* pairs. Along with the type of group we recorded the position registered with a handheld GPS, the number of individuals and the sea state on the Beaufort scale. Flights were suspended when the visibility conditions were not optimal, either because of fog or the sea state exceeded the level 3 of Beaufort scale. The information was introduced into a database developed specifically for this purpose.

The information was clumped into four periods, making each period as similar as possible considering the number of flights. A total of 58 air surveys were clumped in four periods: period 1 from 1999 and 2000 (8 flights) and the other 3 periods comprised from 2004 to 2007 for period 2 (18 flights), period 3 from 2008 to 2012 (17 flights) and period 4 from 2013 to 2016 (15 flights). For each period, the average density per group type was calculated for the entire sampling area.

Later the coast of the surveyed coast was divided into segments of 5 km in length, totalling 124 segments for the 620 km sampled in each flight, where the 0 km is the mouth of the Chubut River (Figure 1). The densities were calculated by dividing the number of whales counted in each segment, weighted by the number of flights performed per period and the area of each segment, calculated as the 5 km segment by 1.5 km (bandwidth), and thus each segment accounted for a 7.5 km<sup>2</sup>. The length of the segment was chosen following Rowntree et al. (2001), who divided the coast into 5 km segments to evaluate the distribution of the Southern Right Whales.

We defined two high densities zones, one inside Golfo Nuevo from Puerto Madryn to Punta Cormoranes and other inside Golfo San José, from Punta Conos to Punta Tehuelche (Figure 1). Thus, we analyzed if there were changes in the distribution of the densities among periods in these zone with Kolmogorov-Smirnov tests (Zar 1999). We also defined the low-density zones as the ones outside the high-density zones (*ie*: Chubut River mouth-Puerto Madryn; Punta Cormoranes-Punta Conos; Punta Tehuelche-Puerto Lobos). Differences in densities among the periods in these zones were assessed by Mann-Whitney U tests (Zar 1999).

#### Results

The overall density of both groups for the whole surveyed area shows for the first period a similar density for *Mother-calf* pairs and *Other* groups. For the second period, the observed increase is similar for both categories (Figure 3). During the third and the fourth period, there is a slight increase in the density of *Mother-calf* pairs, while the *Other* group's density decreases (Figure 3).



Figure 3: Variation of the average density of the sampling area for each established period

Numbering the 5 km segments consecutively from the Chubut river mouth (Figure 1), the coastal areas defined as high densities zones are located between the segment 24 (Puerto Madryn) and segment 49 (Punta Cormoranes) inside Golfo Nuevo and between the segment 90 (Punta Conos) and the segment 101 (Punta Tehuelche) inside Golfo San José (Figure 4).

The highest densities in any of the four periods are observed in the Doradillo (segment 29 and 30) and Playa Fracaso (segment 98 and 101), these two regions are the ones with the highest densities. The maximum mean density was calculated in 3.15 whales per km<sup>2</sup>.



Figure 4: Density of the number of whales per km2 in each segment of 5km performed in the 4 established periods for all the counted whales in the surveyed area without discriminating the group type. *Mother-calf* pairs: white bars, *Other* groups: Black bar, All groups: gray continuous line.

For maximum density zones, Kolmogorov-Smirnov tests were performed, testing if the distribution of densities in each area changed over the four periods. In the zone defined from Puerto Madryn to Punta Comoranes the distribution of the density for the whole population changed from the first period, being stable during the second and third period (Table 1). A greater variation in density was observed during the fourth period, and because of this, there is no difference with the previous years (Figure 5). The described pattern is almost the same for the *Mother-calf* pairs, except that the distribution of the density for this kind of groups is different also for the fourth period (Table 1). This type of group continually increased its density in this zone. The *Other* groups decreased their density in this zone, and this change is evident between the second and the fourth period (Table 1, Figure 5).

In the case of the zone defined between Punta Conos and Punta Tehuelche, the distribution of the densities for all type of groups was more stable across the four periods. For the whole population the only change in the distribution of the density was between

the first and the second period, and the same pattern can be observed with the *Mother-calf* pairs (Table 1). This lack of generalized change in the distribution is related to the large variation of the densities the data show (Figure 5). In the case of the *Other* groups, the distribution shift was between the third and the fourth period (Table 1, Figure 5).

Puerto Madryn - Punta Cormoranes				Punta Conos - Punta Tehuelche			
Period 1	Other Groups			Period 1	Other Groups		
0.068	Period 2			0.152	Period 2		
0.259	0.675	Period 3		0.060	0.999	Period 3	
0.893	0.031	0.259	Period 4	0.905	0.153	0.021	Period 4
Period 1	Mother-calf			Period 1	Mother-calf		
0.005	Period 2			0.153	Period 2		
0.031	0.675	Period 3		0.021	0.905	Period 3	
0.005	0.675	0.259	Period 4	0.334	0.999	0.905	Period 4
Period 1	All Groups			Period 1	All Groups		
0.013	Period 2			0.153	Period 2		
0.031	0.675	Period 3		0.021	0.921	Period 3	
0.068	0.675	0.902	Period 4	0.334	0.617	0.617	Period 4

Table 1: Values of p for the comparison of the distributions of whale densities in the two zones of maximum density for each different group in the 4 periods. In bold and italics are the p that have a significant difference.



Figure 5: Variation of whale densities in the 4 periods for the three groups classified in the two high-density zones.

Figure 6 shows the changes in the densities of the low-density zones. *Mother-calf* pairs for the zone defined from Chubut River Mouth to Puerto Madryn increased their density during the third period ( $U_{2,3}$ : 215; p <0.05). Also, an increase in the density of the *Other* groups from the first to the second period, with a decrease in the density from the second to the third ( $U_{1,2}$ : 205/ $U_{2,3}$ : 540; p< 0.05). In the zone defined from Punta Cormoranes to Punta Conos the *Mother-calf* pairs increased their density only during the third period ( $U_{1vs3}$ : 489/ $U_{2vs3}$ : 437/ $U_{3vs4}$ : 1054/ p <0.05) decreasing afterwards, while for the *Other* 

groups the increase was significant after the first period and remained higher than the first period ( $U_{1vs2}$ : 372/ $U_{1vs3}$ : 519.5/ $U_{1vs4}$ : 511.5/ $U_{2vs4}$ : 994.5; p <0.05). In the area defined from Punta Tehuelche to Puerto Lobos for the *Mother-calf* pairs, there are significant differences between the first period and the rest ( $U_{1vs2}$ : 119/ $U_{1vs3}$ : 109/ $U_{1vs4}$ : 116; p <0.05). The *Other* groups show an increase in density after the first period, but there is also a decrease in density from the second to the fourth period ( $U_{1vs2}$ : 56/ $U_{1vs3}$ : 82/ $U_{1vs4}$ : 71/ $U_{2vs4}$ : 308; p <0.05).



Figure 6: Mean densities of whales in the 4 periods for the three groups classified in the three low-density zones. (the brackets with the asterisk show the periods where there is a significant difference)

The differences observed in the areas of low density show a movement of animals from high areas towards low-density adjacent areas. In the case of the *Mother-calf* pairs for the zones of Chubut River Mount to Puerto Madryn and Punta Cormoranes to Punta Conos there was an increase in the density in the third period and a decrease in the fourth period. Particularly, the density remained higher in the zone from Punta Tehuelche to Puerto Lobos. In the case of the *Other* groups, there was a growth of the density in the second period, in greater measure in Punta Tehuelche to Puerto Lobos, but later the density decreased in the fourth period. This decrease was not compensated by an increase in the high-density zones; contrary to that, the density of others decreased in these zones as well. The changes can also be noted in Figure 4 within a broader context, and the changes in distribution and relocation to low-density areas always is recorded after the global density in the area was near 3 whales per km<sup>2</sup>.

#### Discussion

This is the first study that proposes that a threshold in whale's density in the core areas triggers a density-dependent response. This response includes the movement of Solitary Individuals and Breeding Groups (Other Groups) to adjacent areas when the average density in the area is close to 3 whales per km<sup>2</sup>. This figure must be taken with caution since it is an average from April to December, and the number of whales in the area peak only during August-September (Crespo et al. 2015). During the peak of the season, as much as 15,87 whales per  $\text{km}^2$  can be found in the El Doradillo area but the description of the process can be better viewed using the average. This kind of changes in the distribution of southern right whale in Peninsula Valdés was reported by Rowntree et al. (2001) during the late 80's. Rowntree et al. (2001) propose a movement of whale breeding areas from the outer coast of Peninsula Valdés into the gulfs (Golfo Nuevo and Golfo San José), but no mechanism was proposed. The areas reported as new for the 1990s in Peninsula Valdés are the same as those observed in this work for the first period (Figure 4). Rowntree et al. (2001) considered several factors as gull inflicted wounds or undetected changes in the environment and topography of the area as the possible causation of the observed shift in distribution. Our data lead us to propose that density changes (and hence social causes) can be the main mechanism that promotes the search for new areas and the expansion of the occupied coast in Peninsula Valdés breeding ground, even though this might not have been the same mechanism that drives the changes observed during the late 1980's. Other social interactions are being recorded in the area, besides the already described *Mother-calf* pairs. Solitary individuals engaged in the same activities and cooperative feeding were recorded recently, and hence new interactions never reported before may be shaping the behavior of whales in the area (Argüelles 2017; Arias et al 2017).

*Mother-calf* pairs continue to select areas used in the late 1990s, the so-called highdensity zones (Figure 5). The density in these areas increased, but differentially by type of groups: while *Mother-calf* increased their density, the *Other* groups continue to select this area until an average threshold of 3 whales per km<sup>2</sup> is reached. In every considered period, this is mainly due to the increase in density of *Mother-calf* (the only fraction of the population that is still growing – Crespo et al. (2018)). During the first period, the average annual density reached near 3 whales per km<sup>2</sup> in the area Puerto Madryn- El Doradillo. In the following period a change in the density distribution of whales occurs (Table 1); not only there is an increase in the *Mother-calf* pairs density, but the *Other* groups are more prone to be found in peripheral areas of less density (*eg.* Chubut River Mount-Puerto Madryn) in the second period, as shown in Figure 4. The same pattern can be found in Golfo San José high-density area during the second period. After the mean density approached to 3 whales per km<sup>2</sup>, the density of the *Other* groups increased in the low-density area Punta Tehuelche-Puerto Lobos (Figure 6).

In the fourth period, it is observed that the average annual density of whales in the Doradillo area is close to 3 whales per km<sup>2</sup>. If the same pattern is repeated in the next period, we could hypothesize that an expansion of part of the whales to areas with physical and biological conditions similar to those found in Peninsula Valdes will occur in the next few years, as well as a new increase in the low-density areas.

In a context of a population growth, the expansion into new areas has been recorded, mainly driven by *Solitary Individuals* and *Breeding* groups (Arias et al. 2018). The optimum areas are first occupied by the *Mother-calf* pairs, as the density of this type group increases, the rest of the groups are a move to suboptimal zones (Svendsen 2017). Our results indicate that the recolonization process started at least in the mid-2000s when whales change both, the way they use the habitat related to the type of groups (Crespo et al. 2017) and the areas where they could be found. The expansion of these groups to other areas was observed in the province of Rio Negro with the presence of whales in the area near the San Antonio Bay. In this area, more than 80% of whales are *Solitary Individuals* and *Breeding* Groups (Crespo et al. 2014, Arias et al. 2017). This process has also been recorded in other southern right whale stocks from South Africa, New Zealand and southern Brazil, where *Solitary Individuals* and *Breeding* Groups move to new sites, outside the established breeding area (Barendse and Best 2014, Carroll et al. 2014, Danilewicz et al. 2016).

It is important to monitor the densities in these new areas, as well as the rate of increase in these. The rate of increase of the population is now the combination of the increased recorded in Peninsula Valdés and the growth experienced while recolonizing ancient habitat (Arias et al. 2018, Crespo et al. 2017). Also, it is important to evaluate the habitat suitability of different areas, and to test if the mean density of around 3 individuals per km<sup>2</sup> as a threshold holds outside Peninsula Valdés; and if so, which are the social causes that trigger this density-dependent response.

#### ACKNOWLEDGEMENTS

This paper is dedicated to our colleague Susana N. Pedraza. She was the most influential quantitative researcher at the Marine Mammal Laboratory. This project was supported over the years by Fundación Vida Silvestre Argentina (thanks to Javier Corcuera and Manolo Arias), BIOCON 04 BBVA Foundation, CONICET, ANPCyT, National University of Patagonia, GEF/PNUD 02/018, Southern Spirit SRL (thanks to Tiño Resnik) and Secretaría de Turismo de la Provincia de Chubut. Logistic support was provided by CENPAT and the National University of Patagonia. S. Dans, G. Svendsen, M Degrati, F. Grandi, G. Garaffo, B. Berón Vera, L. Hardtke, F. García, A. Carribero, C. Giesse, D. Valés, R. Loizaga, J. Klaich, S. Leonardi, N. Martínez, V. Milano and C. Durante helped as observers or recorders in the flights. We thank several pilots throughout the period but mainly to Peter Dominguez.

#### References

- Argüelles M.B. 2017. Efectos de la navegación comercial y turística sobre la distribución, abundancia y comportamiento de las ballenas francas (*Eubalaena australis*) en el Golfo Nuevo, Chubut. Ph.D. Thesis. Universidad Nacional del Comahue. 185 p.
- Arias, M., M. A. Coscarella, A. Romero, G. M. Svendsen, M. Ocampo Reinaldo, N. Curcio, E. A. Crespo, and R. González. 2018. Impact of whale-watching on Southern Right Whale (*Eubalaena australis*) in Patagonia: assessing the effects from its beginnings in the context of population growth. Tourism Management Perspectives.
- Arias, M., M. A. Coscarella, G. M. Svendsen, M. A. Romero, N. Curcio, N. Sueyro, E. A. Crespo, and R. González. 2017. Changes in the distribution and abundance of Southern Right Whale Eubalaena australis in San Matías Gulf (Patagonia, Argentina). SC/67a/IWC Southern Right Whale Assessment Workshop.
- Bannister, J. 2001. Status of southern right whales (Eubalaena australis) off Australia. Journal of Cetacean Research and Management **2**:103-110.
- Barendse, J., and P. B. Best. 2014. Shore-based observations of seasonality, movements, and group behavior of southern right whales in a nonnursery area on the South African west coast. Marine Mammal Science **30**:1358-1382.
- Best, P. B., A. Brandão, and D. S. Butterworth. 2001. Demographic parameters of southern right whales off South Africa. J. Cetacean Res. Manage **2**.
- Carroll, E. L., W. J. Rayment, A. M. Alexander, C. S. Baker, N. J. Patenaude, D. Steel, R. Constantine, R. Cole, L. J. Boren, and S. Childerhouse. 2014. Reestablishment of former wintering grounds by New Zealand southern right whales. Marine Mammal Science **30**:206-220.
- Cooke, J. 2012. Southwest Atlantic right whales: updated population assessment from photo-id collected at Península Valdés, Argentina. IWC/64/Rep 1 Annex. IWC Scientific Committee.
- Cooke, J., V. Rowntree, and R. Payne. 2001. Estimates of demographic parameters for southern right whales (*Eubalaena australis*) observed off Península Valdés, Argentina. J. Cetacean Res. Manage:125-132.
- Cooke, J., V. Rowntree, and M. Sironi. 2015. Southwest Atlantic right whales: interim updated population assessment from photo-id collected at Península Valdéz, Argentina. SC/66/IWC Southern Right Whale Assessment Workshop **23**:9 pp.
- Crespo, E. A., S. N. Pedraza, S. L. Dans, M. A. Coscarella, G. M. Svendsen, and M. Degrati. 2011. Number of southern right whales *Eubalaena australis* and population trend in the neighbourhood of Península Valdés during the period 1999-2011 by means of aerial and boat surveys. SC/S11/RW4 Southern Right Whale Assessment Workshop 15.

- Crespo, E. A., S. N. Pedraza, S. L. Dans, M. A. Coscarella, G. M. Svendsen, and M. Degrati. 2014. Number of southern right whales *Eubalaena australis* and population trend in the neighbourhood of Península Valdés during the period 1999-2013 by means of aerial and boat surveys. SC/65b/BRG07 IWC SRW Assessment Workshop.
- Crespo, E. A., S. N. Pedraza, S. L. Dans, M. A. Coscarella, G. M. Svendsen, M. Degrati, J. C. Pedraza, and A. Schiavini. 2015. More whales *Eubalaena australis* and decreasing trend. SC/66/IWC Southern Right Whale Assessment Workshop 20.
- Crespo, E. A., M. A. Coscarella, S. N. Pedraza, S. L. Dans, G. M. Svendsen and M. Degrati. 2017. The Southwestern Atlantic Southern Right Whale, *Eubalaena australis*, still growing but at a decelerated rate. Marine Mammal Science:In Press. SC/67a/IWC Southern Right Whale Assessment Workshop.
- Danilewicz, D., I. B. Moreno, M. Tavares, and F. Sucunza. 2016. Southern right whales (Eubalaena australis) off Torres, Brazil: group characteristics, movements, and insights into the role of the Brazilian-Uruguayan wintering ground. Mammalia.
- Galletti Vernazzani, B., E. Cabrera, and R. L. Brownell. 2014. Eastern South Pacific southern right whale photo-identification catalog reveals behavior and habitat use patterns. Marine Mammal Science **30**:389-398.
- Groch, K., J. Palazzo Jr, P. Flores, F. Adler, and M. Fabian. 2005. Recent rapid increases in the right whale (Eubalaena australis) population off southern Brazil. Latin American Journal of Aquatic Mammals **4**:41-47.
- Hobbs, N. T., and T. A. Hanley. 1990. Habitat evaluation: do use/availability data reflect carrying capacity? The Journal of Wildlife Management:515-522.
- IWC. 2001. Report of the workshop on the comprehensive assessment of right whales: a worldwide comparison. Journal of Cetacean Research and Management **2**:1-60.
- IWC. 2011. Report of the Workshop on the Southern Right Whale Assessment at Buenos Aires, Argentina.
- IWC. 2012. Report of the IWC Workshop on the Assessment of Southern Right Whales. Document SC/64/Rep5 International Whaling Commission. Buenos Aires. 5.
- Payne, R. 1986. Long term behavioral studies of the southern right whale (*Eubalaena australis*). Report of the International Whaling Commission **10**:161-167.
- Payne, R., O. Brazier, E. M. Dorsey, J. S. Perkins, V. Rowntree, and A. Titus. 1981. External features in southern right whales (*Eubalaena australis*) and their use in identifying individuals. Report of the International Whaling Commission.
- Payne, R., V. Rowntree, J. S. Perkins, J. G. Cooke, and K. Lankester. 1990. Population size, trends and reproductive parameters of right whales (*Eubalaena australis*) off Peninsula Valdes, Argentina. Report of the International Whaling Commission:271-278.
- Richards, R. 2009. Past and present distributions of southern right whales (Eubalaena australis). New Zealand Journal of Zoology **36**:447-459.
- Rowntree, V., R. Payne, and D. M. Schell. 2001. Changing patterns of habitat use by southern right whales (*Eubalaena australis*) on their nursery ground at Península Valdés, Argentina, and in their long-range movements. Journal of Cetacean Research and Management **2**:133-143.
- Svendsen, G. M. 2017. Distribución y uso de hábitat de mamíferos marinos en el golfo San Matías.
- Verner, J., M. L. Morrison, and C. J. Ralph. 1986. Wildlife 2000. Modeling habitat relationships of terrestrial vertebrates. University of Wisconsin Press.
- Whitehead, H., R. Payne, and M. Payne. 1986. Population estimate for the right whales off Peninsula Valdes, Argentina, 1971-1976. Report of the International Whaling Commission:71-169.
- Zar, J. H. 1999. Biostatistical analysis. Pearson Education India.