

Report of the Scientific Committee

Virtual Meetings, 27 April-14 May 2021

Annex D: Revised Specifications for Western North Pacific Minke Whales

This report is presented as it was at SC/68C.

There may be further editorial changes (e.g. updated references, tables, figures) made before publication.

**International Whaling Commission
Cambridge, UK, 2021**

Annex D

Specifications for the *In Depth Assessment* of western North Pacific Minke Whales

C. Allison, C.L. de Moor and A.E. Punt

DRAFT – the details of some of these specifications remain to be finalised

A. Basic concepts and stock structure

The objective of this *In Depth Assessment* for western North Pacific minke whales is to review the current status of the stocks and to examine the effect of future catches, for example as set by the Revised Management Procedure (RMP). This assessment has been developed from the *Implementation Simulation Trials* previously used to test the performance of the RMP in scenarios that relate to the actual problem of managing a likely fishery for minke whales in the North Pacific (IWC, 2014b)¹. The trials attempt to bound the range of plausible hypotheses regarding the number of minke whale stocks in the North Pacific, how they feed (by sex, age and month) and recruit and how surveys index them. The underlying dynamics model is age- and sex-structured and allows for multiple stocks.

The region to be managed (the western North Pacific) is divided into 22 sub-areas (see Fig. 1). Future surveys are unlikely to cover sub-areas 1, 2, 3, 4 and 13 (see Table 3) so these sub-areas are taken to be *Residual Areas* in the current trials (although allowance is made for future bycatches from some of these sub-areas – see section D). The term ‘stock’ refers to a group of whales from the same breeding ground.




Fig. 1. The 22 sub-areas used in the *In Depth Assessment* for North Pacific minke whales

¹ Since this *Implementation Assessment* is developed from the *Implementation Simulation Trials* framework, we continue to use the testing nomenclature from the trials (e.g. conditioning rather than fitting).

Three fundamental hypotheses are considered to account for patterns observed in the results from the genetic analyses²:

- (i) there is a single J-stock that occurs to the west of Japan (Sea of Japan and Yellow Sea) and the Pacific coast of Japan (sub-areas 2C, 7CS, 7CN, 11 and 12SW) and a single O-stock in sub-areas to the east and north of Japan (2C, 2R, 3, 4, 7CS, 7CN, 7WR, 7E, 8, 9, 9N, 10E, 11, 12SW, 12NE and 13) (referred to as hypothesis A);
- (ii) as for hypothesis (A), but there is a third stock (Y) that resides in the Yellow sea (sub-areas 1W, 5 and 6W) and overlaps with J-stock in the southern part of sub-area 6W (referred to as hypothesis B); and
- (iii) there are four stocks, referred to Y, J, P, and O, two of which (Y and J) occur in the Sea of Japan, and three of which (J, P, and O) are found to the east of Japan (referred to as hypothesis E). Stock P is a coastal stock.

Sensitivity tests in which there is a C-stock are also conducted based on stock structure hypotheses A and E. The C-stock is found in sub-areas 9 and 9N for the sensitivity test based on stock structure hypothesis A and in these sub-areas as well as sub-area 12NE for the sensitivity test based on stock structure hypothesis E. There is uncertainty regarding whether C-stock is found in sub-area 12NE because of the lack of genetics data for this sub-area.

B. Basic dynamics

Further details of the underlying age-structured model and its parameters can be found in IWC (1991, p.112), except that the model has been extended to take sex-structure into account. The dynamics of the animals in stock j are governed by Equations B.1(a) except for hypothesis E, which allows for dispersal (permanent movement between stocks) as given by Equations B.1(b).

$$N_{t+1,a}^{g,j} = \begin{cases} 0.5 b_{t+1}^j & \text{if } a = 0 \\ (N_{t,a-1}^{g,j} - C_{t,a-1}^{g,j}) \tilde{S}_{a-1} & \text{if } 1 \leq a < x \\ (N_{t,x}^{g,j} - C_{t,x}^{g,j}) \tilde{S}_x + (N_{t,x-1}^{g,j} - C_{t,x-1}^{g,j}) \tilde{S}_{x-1} & \text{if } a = x \end{cases} \quad (\text{B.1a})$$

$$N_{t+1,a}^{g,j} = \begin{cases} 0.5 b_{t+1}^j & \text{if } a = 0 \\ \sum_{j \neq j'} [(1 - D^{j,j'}) (N_{t,a-1}^{g,j} - C_{t,a-1}^{g,j}) \tilde{S}_{a-1} + D^{j,j'} (N_{t,a-1}^{g,j'} - C_{t,a-1}^{g,j'}) \tilde{S}_{a-1}] & \text{if } 1 \leq a < x \\ \sum_{j \neq j'} [(1 - D^{j,j'}) ((N_{t,x}^{g,j} - C_{t,x}^{g,j}) \tilde{S}_x + (N_{t,x-1}^{g,j} - C_{t,x-1}^{g,j}) \tilde{S}_{x-1}) \\ + D^{j,j'} ((N_{t,x}^{g,j'} - C_{t,x}^{g,j'}) \tilde{S}_x + (N_{t,x-1}^{g,j'} - C_{t,x-1}^{g,j'}) \tilde{S}_{x-1})] & \text{if } a = x \end{cases} \quad (\text{B.1b})$$

where $N_{t,a}^{g,j}$ is the number of animals of gender g and age a in stock j at the start of year t ;
 $C_{t,a}^{g,j}$ is the catch (in number) of animals of gender g and age a in stock j during year t (whaling is assumed to take place in a pulse at the start of each year);
 b_t^j is the number of calves born to females from stock j at the start of year t ;
 \tilde{S}_a is the survival rate = e^{-M_a} where M_a is the instantaneous rate of natural mortality (assumed to be independent of stock and sex); and
 x is the maximum age (treated as a plus-group); and
 $D^{j,j'}$ is the dispersal rate (i.e. the probability of an animal moving permanently) from stock j to j' . It is assumed that the numbers dispersing from the j -stock to the j' -stock are the same as from the j' -stock to the j -stock at unexploited equilibrium and that the proportion of calves dispersing from the j -stock to the j' -stock at equilibrium is the same as that from the j' -stock to the j -stock.

Note that projections start in year $t=2021$.

For computational ease, the numbers-at-age by sex are updated at the end of each year only, even though catching is assumed to occur from March to October. This simplification is unlikely to affect the results substantially for two reasons: (1) catches are at most only a few percent of the number of animals selected to the fisheries; and (2) sightings survey estimates are subject to high variability so that the resultant slight positive bias in abundance estimates is almost certainly inconsequential.

² See IWC, 2020 pp376-381 for details of the data and analyses used in the development of these hypotheses.

C. Births

Density-dependence is assumed to act on the female component of the mature population. The convention of referring to the mature population is used here, although this actually refers to animals that have reached the age of first parturition.

$$B_t^j = B^j N_t^{f,j} \{1 + A^j (1 - (N_t^{f,j} / K^{f,j})^{z^j})\} \quad (\text{C.1})$$

where B^j is the average number of births (of both sexes) per year for a mature female in stock j in the pristine population;

A^j is the resilience parameter for stock j ;

z^j is the degree of compensation for stock j ;

$N_t^{f,j}$ is the number of 'mature' females in stock j at the start of year t :

$$N_t^{f,j} = \sum_{a=a_m}^x N_{t,a}^{f,j} \quad (\text{C.2})$$

a_m is the age-at-first-parturition; and

$K^{f,j}$ is the number of mature females in stock j in the pristine (pre-exploitation, written as $t=-\infty$) population:

$$K^{f,j} = \sum_{a=a_m}^x N_{-\infty,a}^{f,j} \quad (\text{C.3})$$

The values of the parameters A^j and z^j for each stock are calculated from the values for $MSYL^j$ and $MSYR^j$ (Punt, 1999). Their calculation assumes harvesting equal proportions of males and females.

D. Catches

The operating model considers two sources for non-natural mortality: direct catches and bycatches (which are also referred to as incidental catches). In future ($t > 2020$), the former are set externally (e.g. by the RMP or specified as a time-series of fixed removals by sub-area), while the latter are a function of abundance and future fishery effort.

In cases in which the total catch limit (e.g. as set by the RMP) is less than the level of incidental catch, the total removals are taken to be the incidental catch only whereas if this total catch limit exceeds the incidental catch (if any), the level of the commercial removals is taken to be the difference between the total catch limit and the best estimate of the incidental catch (see 'Future incidental catches' below).

D.1 Direct catches

The direct historical (pre-2021) catch series used are listed in Appendix 1 and include both commercial and special permit catches. Details of the sources of the catch data are given in Allison (2011). The baseline trials use the 'best' direct catch series, and an alternative 'high' catch series is used in Trial 4. Trials 8 and 9 test the effect of the method used to allocate historical catches between sub-areas 5 and 6W. If catch limits are set by the RMP, it will use the 'best' series in all cases; i.e. it will use what are in effect incorrect catches for Trials 4, 8 and 9 to examine the implications of uncertainty about historical catches. Catch limits are set by *Small Area*. (Catches are always reported by *Small Area*).

Catches and bycatches are removed month by month from each sub-area. It is assumed that whales are homogeneously distributed across a sub-area (excepting in sub-areas 7CS and 7CN in the future), so historical catches and the future catch limits for a sub-area are allocated to stocks by sex and age relative to their true density within that sub-area, and a catch mixing matrix V that depends on sex, age and time of the year (and may also depend on year), i.e.

$$C_{t,a}^{g,j} = \sum_k \sum_q F_t^{g,k,q} V_{t,a}^{g,j,k,q} S_a^g \tilde{N}_{t,q,a}^{g,j} \quad (\text{D.1})$$

$$F_t^{g,k,q} = \frac{C_t^{g,k,q}}{\sum_{j'} \sum_{a'} V_{t,a'}^{g,j',k,q} S_{a'}^g \tilde{N}_{t,q,a'}^{g,j'}} \quad (\text{D.2})$$

where $F_t^{g,k,q}$ is the exploitation rate in sub-area k on fully recruited ($S_a^g \rightarrow 1$) animals of gender g during month q of year t ;

S_a^g is the selectivity on animals of gender g and age a :

$$S_a^g = (1 + e^{-(a - a_{s0}^g) / \delta^g})^{-1} \quad (\text{D.3})$$

$\tilde{N}_{t,q,a}^{g,j}$ is the number of animals of gender g and age a in stock j at the start of month q in year t after removal of catches in earlier months and after removal of any bycatches in month q ;

- a_{50}^g, δ^g are the parameters of the (logistic) selectivity ogive for gender g ; and
 $C_t^{g,k,q}$ is the catch of animals of gender g in sub-area k during month q of year t (see Appendix 1 for the historical catches).

Each entry in the catch mixing matrix, $V_{t,a}^{g,j,k,q}$, is the fraction of males/females of age a from stock j that are found in sub-area k during month q of year t . The catch mixing matrix is different for each month to reflect the effects of migration between the breeding and the feeding grounds and back. Appendix 2 lists the catch mixing matrices considered. The matrices are based on the presence/absence matrices developed at the First Intersessional Workshop (see IWC, 2020) and represent the relative fraction of an age-class in each of the sub-areas during the months March–October. Once the values of the parameters related to mixing rates (the γ s – see section F) are specified (these are estimated separately for each trial and each replicate during the conditioning process), the catch mixing matrices can be converted to fractions of each age-class in each sub-area. The values for the γ parameters are selected to mimic available data (see Section F).

Catch mixing matrices are specified for ages 4 and 10 (these being three years below and above the assumed age-at-50%-maturity). Few animals of age 4 are mature while most of age 10 are. The catch mixing matrices for ages 0–3 are assumed to be the same as that for age 4, and those for ages 11+ the same as that for age 10. The catch mixing matrices for ages 5–9 are set by interpolating linearly between those for ages 4 and 10.

The trials model whale movements in the eight-months from March to October. In order to account for historical direct and incidental catches outside these months, all catches in January–March are modelled as being taken in March and the catches after October are assumed to have been taken in October. The historical direct catches by sex, sub-area, month and year are given in Appendix 1.

The trials are conducted assuming that the sub-areas for which future catch limits might be set are:

sub-area	7CS and 7CN	April to October (coastal/pelagic whaling outside a specified distance ³)
	7WR and 7E	April to October (pelagic whaling)
	8 and 9	April to October (pelagic whaling)
	11	April to October (coastal and pelagic whaling)
	12	April to October (coastal and pelagic whaling)

Future ($t > 2020$) commercial catches are allocated to sex, sub-area, month and year using the equation:

$$C_t^{g,k,q} = C_t^k Q_t^{g,k,q} \quad (\text{D.4})$$

$Q_t^{g,k,q}$ is the fraction of the commercial catch in sub-area k of gender g that is taken during month q , the values of which are given in Table 1a; and

C_t^k is the commercial catch limit for sub-area k and year t ($t > 2020$). Note that C_t^k is equal to the total catch limit (e.g. as set by the RMP) less any reported incidental catch (constrained to be non-negative).

Entries in the Q matrix are determined by the options related to the sub-areas for which catch limits might be set; the non-zero entries (see Table 1a) reflect the historical breakdown of catches over the last 10 years of commercial whaling (1978–87) within each sub-area. In sub-areas for which there was no catch between 1978–87 (7E, 8 and 9), the entries in the Q matrix are set using the entire historical commercial and scientific catch in these sub-areas. In some instances where regulations limited the commercial whaling season, the matrix entries have been adjusted using the special permit data.

Future commercial catches are allocated to stock as described above (Equations D.1 and D.2) except in sub-areas 7CS and 7CN where the genetic data show differences between nearshore and offshore catches. It is assumed future catches will be taken offshore and are allocated to stock based on the mixing proportions set using genetic data from special permit samples only (Table 2a). The process of allocating removals to stock within sub-areas 7CS and 7CN involves first denoting the modelled mixing proportion used when conditioning, $R^{k,q}$, as:

$$R^{k,q} = \sum_{t=1996}^{2016} P_{1+,t}^{J/JE,k,q} \left/ \sum_j \sum_{t=1996}^{2016} P_{1+,t}^{j,k,q} \right.$$

where $P_{1+,t}^{j,k,q}$ is the average number of 1+ animals from stock j in sub-area k in month q of year t .

³Operations preliminarily being considered would be limited ‘to outside a certain distance from the coast to minimise catch of J-stock whales’ (see IWC, 2020 p387). The 2013 trials were conducted assuming whaling would be outside 10 n.miles.

The mixing proportions obtained from the offshore samples, $\tilde{R}^{k,q}$, are given in Table 2a. The proportion of J-stock animals in some future year would normally be $P_{1+,t}^{J,k,q} / (P_{1+,t}^{J,k,q} + P_{1+,t}^{P,k,q} + P_{1+,t}^{O,k,q})$. For sub-areas 7CS and 7CN in future this equation is adjusted to:

$$(\tilde{R}^{k,q} \neq R^{k,q}) : \alpha^{k,q} P_{1+,t}^{J,k,q} / (\alpha^{k,q} P_{1+,t}^{J,k,q} + P_{1+,t}^{P,k,q} + P_{1+,t}^{O,k,q}) \text{ where } \alpha^{k,q} = \frac{(1 - R^{k,q})\tilde{R}^{k,q}}{(1 - \tilde{R}^{k,q})R^{k,q}} \quad (\text{D.4a})$$

The $\alpha^{k,q}$ factor is then applied to the recruited population from J-stock in sub-area k and month q when setting the commercial catch by stock using Equations D.1 and D.2.

Table 1a

The Q matrix used to allocate future commercial catches for a sub-area to sex and month. The entries give the percentage of the catch in sub-area k that is taken by sex and month for sub-areas other than *Residual Areas*. Dashes indicate sub-areas/months for which catch limits are defined to be zero. See text for description of how the entries are set. Values are set using catches taken up to and including 2018.

Sub-area	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct
Males																
7CS	-	24.3	21.5	10.1	4.8	0.8	0.3	-	-	21.7	12.6	2.8	0.7	0.3	-	-
7CN	-	-	0.8	8.2	15.5	15.3	23.9	11.9	-	0.1	0.4	4.9	6.9	3.5	5.3	3.1
7WR	-	0.9	45.0	30.3	2.8	0.9	6.4	-	-	-	8.3	2.8	2.8	-	-	-
7E	-	-	32.9	19.3	1.9	7.2	12.6	1.0	-	-	3.9	1.9	5.3	5.3	8.7	-
8	-	-	12.8	33.6	31.9	4.4	3.0	2.0	-	-	2.7	2.0	3.4	2.0	0.7	1.7
9	-	-	5.4	13.6	30.4	36.3	2.9	-	-	-	1.5	1.8	2.7	4.9	0.5	-
11	-	1.3	5.5	9.6	9.6	4.0	3.0	0.6	0.1	10.6	19.3	18.5	10.7	4.5	2.3	0.4

Table 1b

QB matrix: the percentage of the incidental catch in sub-area k that is taken by sex and month. The values are set using all available bycatches known by sub-area, sex and month, up to and including 2016 (Japan) and 2017 (Korea). There are no known incidental catches in other sub-areas.

Sub-area	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Mar	Apr	May	Jun	Jul	Aug	Sept	Oct	Sample size
Males																	
1E	17.1	9.21	1.32	9.21	1.32	0	0	3.95	18.4	6.58	10.5	7.89	6.58	2.63	0	5.26	76
2C	15.1	4.3	2.42	0.81	1.08	0.54	0	14.2	24.7	1.88	3.76	2.42	2.69	1.61	0.27	24.2	372
5	5.17	3.45	10.3	19.8	1.72	2.59	1.72	12.1	9.48	4.31	7.76	7.76	3.45	0	1.72	8.62	116
6W	13.3	5.91	6.6	4.75	2.67	3.01	4.17	14.6	13.2	4.98	4.63	6.14	1.16	1.51	1.74	11.6	863
6E	15.5	9.88	6.79	2	2.5	2.5	1.2	9.08	16.7	9.28	6.29	2.69	1.7	2.1	1.1	10.8	1002
7CS	7.89	5.02	10.4	7.17	2.51	1.08	0.36	11.5	10	8.96	9.32	8.6	2.15	1.43	1.08	12.5	279
7CN	4.19	4.79	3.59	8.38	7.19	1.8	1.2	9.58	2.99	8.98	12	9.58	6.59	2.99	1.8	14.4	167
10E	0	0	0	0	0	5.56	0	55.6	0	0	0	5.56	0	0	0	33.3	18
11	0	0	0	4.08	0	0	6.12	24.5	0	0	18.4	18.4	4.08	0	2.04	22.4	49

Table 2a

Time-invariant fixed proportions by stock to be used in removing **future commercial catches** from sub-areas 7CS and 7CN for each for stock hypothesis, based on the number of sampled whales that were assigned to each stock using the genetic data⁴ limited to special permit samples only [in the 2013 trials this was limited to >10nm]. The values are set using data from 1996-2016.

Hypothesis	Sub-Area	Months	Sample size		Proportion	
			J-Stock	O-Stock	J-Stock	O-Stock
A & B	7CS	Apr	48	138	0.258	0.742
A & B	7CS	May	89	255	0.259	0.741
A & B	7CS	Jun-Sep	4	75	0.051	0.949
A & B	7CN	Apr-Jun	12	139	0.079	0.921
A & B	7CN	Jul-Dec	169	645	0.208	0.792

Hypothesis	Sub-Area	Months	Sample size			Proportion		
			J-Stock	P-Stock	O-Stock	J-Stock	P-Stock	O-Stock
E	7CS	Apr	0	188	0	0.000	1.000	0.000
E	7CS	May	0	303	24	0.000	0.927	0.073
E	7CS	Jun-Sep	0	5	73	0.000	0.064	0.936
E	7CN	Apr-Jun	2	28	109	0.014	0.201	0.784
E	7CN	Jul-Dec	10	574	225	0.012	0.710	0.278

⁴ From the data file "Data_NPM_190226_v3.csv", based on "stock90" for Hypotheses A&B and "geneland.stock2" for Hypothesis E, using special permit data only. The months are based on the same month-split used in 2013 for commercial catches. There were no special permit catches in sub-areas 7CN & 7CS in Jan-Mar or in sub-area 7CS in Oct-Dec.

Table 2b

Time-invariant fixed proportions by stock to be used in removing **bycatch** from sub-areas 7CS and 7CN for each for stock hypothesis, based on the number of sampled whales that were assigned to each stock using genetic data⁵ limited to bycatch only, using data from 2001-2016.

Hypothesis	Sub-Area	Months	Sample size		Proportion	
			J-Stock	O-Stock	J-Stock	O-Stock
A & B	7CS	Jan-Apr	43	34	0.558	0.442
A & B	7CS	May	16	31	0.340	0.660
A & B	7CS	Jun-Dec	86	34	0.717	0.283
A & B	7CN	Jan-Jun	38	44	0.463	0.537
A & B	7CN	Jul-Dec	51	15	0.773	0.227

Hypothesis	Sub-Area	Months	Sample size			Proportion		
			J-Stock	P-Stock	O-Stock	J-Stock	P-Stock	O-Stock
E	7CS	Jan-Apr	0	73	1	0.000	0.986	0.014
E	7CS	May	0	49	2	0.000	0.961	0.039
E	7CS	Jun-Dec	0	118	1	0.000	0.992	0.008
E	7CN	Jan-Jun	12	69	0	0.148	0.852	0.000
E	7CN	Jul-Dec	13	59	0	0.181	0.819	0.000

D.2 Incidental catches (also known as bycatches)

Incidental catches of minke whales are known to occur off Japan (in sub-areas 1E, 2C, 6E, 7CS, 7CN, 10E and 11 and small numbers in 6W) and the Republic of Korea (sub-areas 5 and 6W and small numbers in 1W).

Japan: It has been obligatory to report bycatches in Japan since 2001 since when the bycatch numbers are considered to be reliable. Earlier bycatches are believed to be under-reported based on the sudden increase in reported bycatches in 2001. In view of this, the relationship between bycatch and set-net effort is integrated into the conditioning process, with the advantage that the method is independent of the reporting rate prior to 2001. The reporting rate since 2001 is assumed to be constant at 100% (except in Trial 4 – see below).

Almost all of the reported bycatch off Japan occurred in set-net fisheries. Three types of set nets are used off Japan: large-scale (excluding salmon nets), salmon nets and small-scale. For fishing gears other than set-nets, incidental catch, retention and marketing of whales are prohibited by the 2001 regulation and a diagnostic DNA registry is used to deter illegal distribution of whales caught. Ideally, the catch by each gear type should be modelled separately to allow the historical (pre-2001) bycatch to be predicted. However, information on numbers of catches by net type is not available. Therefore, in the 2013 Implementation, the historical bycatches for each sub-area were set using the total number of incidental catches and the combined number of large-scale and salmon nets in each sub-area. The numbers of salmon nets since 2006 are not available and as the numbers caught in salmon nets are small in comparison to those from large-scale nets (see Appendix 1). In the current trials, the historical bycatches are extrapolated using the total number of incidental catches and the number of large-scale nets only in each sub-area over the period 2002-2018. For the best effort series, the number of nets from Japan is extrapolated from 1946 to 1969 assuming a linear relationship from 0 in 1935 to the known number in 1970 (Tobayama *et al.*, 1992). Incidental catches before 1946 are ignored because although some set-nets were in operation before 1946 (Brownell, pers. comm.) the numbers are highly uncertain and are sufficiently small that they are unlikely to affect the conditioning process.

The year 2001 is excluded from the fitting because the catch data are incomplete, as the new regulations date from June 2001. A high effort series is also generated, for use in Trial 4, in which the number of nets is double the best-case values from 1946-1969, up to a maximum equal to the number of nets in 1969. In Trial 4 all bycatches are assumed to be under-reported and are adjusted upward by a factor of 2.

Korea. The same method is used as for Japan above except the incidental catch numbers from 1996-2009 (sub-area 6W) and 2000-2009 (sub-area 5) are used to extrapolate backwards and the incidental catch numbers are adjusted to allow for underreporting. The bycatches in sub-area 6W (the East Sea) are adjusted upward by a factor of 2. The factor 2 is based on DNA profiling and a capture-recapture analysis of market products that estimated a total of 887 whales going through Korean markets from 1999-2003, in comparison to the reported catch of 458 whales (Baker *et al.*, 2007). The baseline trials assume that the bycatches in the Yellow Sea (sub-area 5) are fully reported as there is no evidence of under-reporting. The ‘high’ effort series for sub-area 5 used in Trial 4 will apply the same estimate of under-reporting as for sub-area 6W (i.e. a factor of 2) and the number of nets is set to the maximum of either double the base-case values or the number of nets in 1969.

To account for bycatch prior to 1996, the average for the *adjusted* takes are used to extrapolate backwards to 1946 based on fisheries effort using the same approach as for Japan. Incidental catches before 1946 are ignored as for Japan.

⁵ From the data file “Data_NPM_190226_v3.csv”, based on “stock90” for Hypotheses A&B and “geneland.stock2” for stock hypothesis E, using bycatch data only. The months are based on the same month-split used in 2013 for bycatches.

China. There are no data on incidental catches off China, although they are known to occur. The trials therefore consider two (essentially arbitrary) scenarios: (i) the incidental catch by China is twice that reported by Korea in sub-area 5; and (ii) incidental catches off China are ignored. The first of the options forms part of the baseline specifications and the second is included in a sensitivity test (Trial 12) to determine the effects of the base case assumptions.

Allocation to sex and month. Bycatches by sex, sub-area, month and year, $C_{B,t}^{g,k,q}$, are set using the equation:

$$C_{B,t}^{g,k,q} = C_{B,t}^k Q_B^{g,k,q} \quad (D.5)$$

$Q_B^{g,k,q}$ is the fraction of the bycatch of gender g in sub-area k which is taken during month q and, the values of which are given in Table 1b; and

$C_{B,t}^k$ is the bycatch in sub-area k and year t (as estimated by the model).

To avoid a proliferation of sub-areas and to avoid the need for finer time-steps than month, incidental catches in sub-areas other than 7CS and 7CN are apportioned to stock and age class in the same way as for the commercial catches in Equations D.1 and D.2, but assuming that the bycatch is taken uniformly from all age classes (i.e. selectivity=1). Thus

$$C_{B,t}^{g,j} = \sum_k \sum_q F_{B,t}^{g,k,q} V_{t,a}^{g,j,k,q} \ddot{N}_{t,q,a}^{g,j}$$

$F_{B,t}^{g,k,q}$ is the bycatch removal rate for gender g in sub-area k (all sub-areas except 7CS and 7CN) during month q of year t

$$F_{B,t}^{g,k,q} = \frac{C_{B,t}^{g,k,q}}{\sum_{j'} \sum_{a'} V_{t,a'}^{g,j',k,q} \ddot{N}_{t,q,a'}^{g,j'}}$$

In sub-areas 7CS and 7CN, (where the genetic data show differences between nearshore and offshore catches) bycatches are taken nearshore and so are allocated to stock using mixing proportions calculated from the number of sampled whales that were assigned to each stock using genetic data from bycatches only (Table 2b).

$$\tilde{N}_{t,q,a}^{g,j} = \ddot{N}_{t,q,a}^{g,j} (1 - V_{t,a}^{g,j,k,q} F_{B,t}^{g,k,q}) \quad \text{for all sub-areas except 7CS and 7CN and}$$

$$\tilde{N}_{t,q,a}^{g,j} = \ddot{N}_{t,q,a}^{g,j} (1 - F_{B,t}^{g,k,q,j}) \quad \text{for sub-areas 7CS and 7CN,}$$

$F_{B,t}^{g,k,q,j}$ is the removal rate due to bycatch of gender g and stock j in sub-area k (sub-areas 7CS and 7CN) during month q of year t .

$$F_{B,t}^{g,k,q,j} = \frac{p_B^{k,q,j} C_{B,t}^{g,k,q}}{\sum_{a'} \ddot{N}_{t,q,a}^{g,j}} \quad \text{where } p_B^{k,q,j} \text{ is given by Table 2b; and}$$

$C_{B,t}^{g,k,q}$ is the bycatch of animals of gender g in sub-area k during month q of year t (given by Equation D.5).

The historical bycatch model: The historical bycatch $C_{B,t}^k$ in sub-area k in year t is given by:

$$C_{B,t}^k = A^k P_t^k E_t^k \quad (D.6)$$

where A^k is the bycatch constant, E_t^k is the number of nets in sub-area k during year t and P_t^k is the total population size (including calves) in sub-area k in year t averaged over all 8 time periods. In Trial 17, the abundance P_t^k in Equation D.6 is replaced by $\sqrt{P_t^k}$ to test an alternative assumption for the relationship between bycatch and abundance and the impact of possible saturation effects. The values of the bycatch constants are set by fitting during the conditioning process (see section F). In years where actual numbers of bycatches are known, these are the values removed from the population rather than the model estimated values.

The recent bycatches and the numbers of set-nets by type, year and area are listed in Appendix 1. Further details are given in Annex H of IWC (2012a).

Future bycatches: Future bycatches by sub-area (except in sub-areas 7CS and 7CN) are generated assuming that the exploitation rate due to bycatch in the future equals that estimated for the trial in question for the most recent five-years of data used in the conditioning process, i.e.:

$$C_{B,t}^k = \bar{F}^k P_t^k \quad (\text{D.7})$$

where $C_{B,t}^k$ is the bycatch in sub-area k in year t , P_t^k is the total population (including calves) in sub-area k during year t averaged over all 8 time periods (March–October), and \bar{F}^k is the average exploitation rate (sum over years of the known bycatch divided by the sum over years of P_t^k) over the last five years of the period used for conditioning (2016–20 for sub-areas off Japan and 2015–19 for those off Korea), i.e. F is reset for each of the 100 simulations within a trial. Thus, the future bycatch by sex, month and sub-area is given by:

$$C_{B,t}^{g,k,q} = Q_B^{g,k,q} \bar{F}^k P_t^k \quad (\text{D.7a})$$

For Trial 17, the abundance P_t^k in Equation D.7a is replaced by $\sqrt{(P_t^k)}$.

To avoid possible dis-proportionate bycatches of J- to O-stock whales, Equation (D.7a) is replaced with (D.7b) in sub-areas 7CS and 7CN.

$$C_{B,t}^{g,k,q} = \tilde{P}_t^k \bar{F}^k Q_B^{g,k,q} \quad (\text{D.7b})$$

where $\tilde{P}_t^{k,q}$ is the availability-weighted population size in sub-area k during month q :

$$\tilde{P}_t^{k,q} = (P_t^{k,q,J} + \lambda^{k,q} P_t^{k,q,O}) \frac{\bar{P}^{k,q,J} + \bar{P}^{k,q,O}}{\bar{P}^{k,q,J} + \lambda^{k,q} \bar{P}^{k,q,O}} \quad (\text{D.8})$$

where $\bar{P}^{k,q,j}$ is the average number (including calves) of stock j animals in sub-area k during month q over the last five years of the period used for conditioning;

$P_t^{k,q,j}$ is the total population size (including calves) of stock j in sub-area k during month q of year t ;

$\lambda^{k,q}$ is a relative availability factor for J whales relative to O whales:

$$\lambda^{k,q} = \frac{(1 - \ddot{P}^{k,q})}{\ddot{P}^{k,q}} \frac{\bar{P}^{k,q,J}}{\bar{P}^{k,q,O}} \quad (\text{D.9})$$

$\ddot{P}^{k,q}$ is the weighted mean proportion of J-stock in sub-area k during month q (as given in Table 2b).

This bycatch is allocated to stock as follows:

$$C_{B,t}^{g,k,q,J} = \frac{P_t^{g,k,q,J}}{\lambda^{k,q} P_t^{g,k,q,O} + P_t^{g,k,q,J}} C_{B,t}^{g,k,q} \quad (\text{D.10a})$$

$$C_{B,t}^{g,k,q,O} = \frac{\lambda^{k,q} P_t^{g,k,q,O}}{\lambda^{k,q} P_t^{g,k,q,O} + P_t^{g,k,q,J}} C_{B,t}^{g,k,q} \quad (\text{D.10b})$$

where $P_t^{g,k,q,j}$ is the total population size (including calves) of animals of gender g from stock j in sub-area k during month q of year t .

Reported bycatches

A single series of historical bycatches will be used for all of the trials when applying the RMP (i.e. for calculating catch limits), irrespective of the true values of the bycatches, which differ both among trials and simulations within trials. The estimate of the historical bycatches used by the CLA will be set to the averages of the predicted bycatches based on the fit to the actual data⁶ of the operating model for the six baseline trials (i.e. using the ‘best fit’ simulation (0)). The series will be generated after conditioning is complete (see Appendix 1).

The future bycatches used when applying the RMP are the true bycatches in all sub-areas⁷, except for Trial 4 (in which the estimated bycatches are in error to reflect the under-estimation of bycatch inherent in these trials) and Trial 12 (in which the bycatch by China is taken to be zero).

⁶ In the case of sub-area 6W the actual data is the *adjusted* bycatch data.

⁷ Including sub-area 6W since the best estimate of bycatches in this area is the adjusted figure.

E. Generation of data

In 2013, the *Implementation Simulation Trials* (IWC, 2014b) used to test the performance of the RMP required estimates of future abundance to be generated. This is retained in the control program and so is documented below, although it is unnecessary for the current assessment.

The plan for future sightings surveys is listed in Table 3a. Surveys will be conducted by Japan in sub-areas 7CS, 7CN, 7WR, 7E, 8, 9, 11, 12SW and 12N. Additional surveys will be conducted by Japan in sub-areas 6E, 10W, 10E and by Korea in sub-areas 5 and 6W (see IWC, 2020 p382), but they are not listed here as they are not required for setting future catch limits and so are not modelled in the trials. Table 3b shows how surveys will be combined for areas that are combinations of sub-areas.

The estimates of absolute abundance (and their associated CVs) that are provided to the CLA for the years prior to management are given in Table 4. Estimates of abundance are generated for any surveys which have already been conducted but for which the results are not yet available.

The sightings mixing matrix for a year in which a survey takes place is the average of the catch mixing matrices over the two survey months in that year (April-May for surveys to the west of Japan or August-September for the remainder). The values for the parameters of the various distributions have been selected to achieve CVs for *Small Areas* comparable to those for the surveys in Table 6. The future estimates of abundance for a *Small Area* (say *Small Area E*) are generated using the formula:

$$\hat{P} = PYw / \mu = P^* \beta^2 Yw \quad (\text{E.1})$$

Y is a lognormal random variable $Y = e^\varepsilon$ where $\varepsilon \sim N[0, \sigma^2]$ and $\sigma^2 = \ln(\alpha^2 + 1)$;

w is Poisson random variable with $E(w) = \text{var}(w) = \mu = (P / P^*) / \beta^2$; (Y and w are independent);

P is the average current total (1+) population size in the *Small Area* (E) over the survey period:

$$P = P_t^E = \frac{1}{2} \sum_{k \in F} \sum_{q \in \text{SurveyPeriod}} \sum_j \sum_g \sum_{a=1}^x (V_{t,a}^{g,j,k,q} N_{t,a}^{g,j}) \quad (\text{E.2})$$

P^* is the reference population level, and is equal to the mean total (1+) population size in the *Small Area* prior to the commencement of exploitation in the area being surveyed; and

F is the set of sub-areas making up *Small Area E*.

Note that under the approximation $CV^2(ab) \approx CV^2(a) + CV^2(b)$: $E(\hat{P}) \approx P$ and $CV^2(\hat{P}) \approx \alpha^2 + \beta^2 P^* / P$

For consistency with the first stage screening trials for a single stock (IWC, 1991, p.109; 1994, pp.85-86), the ratio $\alpha^2 : \beta^2 = 0.12 : 0.025$, so that:

$$CV(\hat{P}) = \tau(0.12 + 0.025 P^* / P)^{1/2} \quad (\text{E.3})$$

and the CV of a survey estimate prior to the commencement of exploitation in the area being surveyed would be:

$$\sqrt{(\alpha^2 + \beta^2)} = 0.38\tau \quad (\text{E.4})$$

The values of τ applicable to each sub-area are calculated separately for each replicate once the conditioning has been accomplished by substituting the true value of the CV for each abundance estimate used in conditioning (Table 6)⁸ and the corresponding model depletion level into Equation E.3. If more than one abundance estimate exists for a particular sub-area, the value assumed for τ is calculated taking the true CV to be the root mean square of the values obtained from the abundance estimates for that sub-area, and the depletion to be the mean value over the corresponding years.

An estimate of the CV, X_t , is also generated for each sightings estimate, \hat{P}_t :

$$X_t = \sqrt{(\sigma_t^2 \chi^2 / n)} \quad (\text{E.5})$$

where $\sigma_t^2 = \ln(1 + \alpha^2 + \beta^2 P^* / \hat{P}_t)$, and χ^2 is a random number from a Chi-square distribution with $n=10$ degrees of freedom. The value 10 is chosen to roughly indicate the number of trackline segments in a sightings survey in a *Small Area*.

The trials will be based on the use of two alternative values for $g(0)$ in the conditioning process: $g(0) = 0.798$ (the base case value) and $g(0)=1$ (Trial 3) (IWC, 2012a, p.417; Okamura *et al.*, 2010). When $g(0) = 0.798$ the values of the operating model abundances are multiplied by this factor when setting the future survey estimates of abundance.

⁸ Excluding zero, minimum and maximum estimates and those assumed to apply to adjacent areas.

Table 3a

Past and planned future Japanese surveys to the North and East of Japan. The survey coverage is given in parentheses. Future coverage in sub-areas 7CN, 7WR and 7E is expected to be similar to the values below (because of territorial issues). Coverage in sub-areas 8 and 9 assumes that future surveys include the Russian EEZ. Future coverage in sub-areas 11 and 12SW (of 30.1% and 48.9% respectively) excludes areas in the Russian EEZ that cannot be surveyed until the resolution of territorial issues with Japan. Future coverage in sub-area 12NE (of 46.4) reflects the area which cannot be surveyed in the North and East because of Russian restrictions. * Estimate=0; # surveys covered different parts of sub-area 12NE each year.

	7CS	7CN	7WR	7E	8	9	11	12SW	12NE
1990	-	-	-	-	Aug (62%)	Aug (61%)	Aug-Sep(100%)	Aug-Sep(100%)	Aug-Sep(100%)
1991	Aug-Sep*(100%)	Aug-Sep(100%)	Aug-Sep(100%)	-	-	-	-	-	-
1992	-	-	-	-	-	-	-	-	Aug-Sep (100%)
1999	-	-	-	-	-	-	Aug-Sep(100%)	-	Aug-Sep (64%)
2000	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-
2002	-	-	-	-	Jun-Jul*(65%)	-	-	-	-
2003	-	May-Jun (27%)	-	-	-	Jul-Sep (33%)	Aug-Sep (34%)	Aug-Sep(100%)	Aug-Sep (41%)
2004	May (37%)	May-Jun (89%)	My-Jun (57%)	Jun (41%)	-	-	-	-	-
2005	-	-	-	-	May-Jul (65%)	-	-	-	-
2006	Jun-Jul (100%)	-	-	My-Jun (57%)	May-Jul (65%)	-	-	-	-
2007	-	-	Jun-Jul (89%)	Jun-Jul* (57%)*	Jun-Jul (65%)	-	Aug-Sep (20%)	-	-
2008	Aug* (100%)	Aug* (75%)	Aug* (89%)	Aug* (57%)	Jul-Aug*(65%)	Jul-Aug (87%)	-	-	-
2009	May-Jun (100%)	May-Jun (75%)	May-Jun (89%)	May-Jun (57%)	May-Jun (65%)	May-Jun (63%)	-	-	-
2010	-	-	-	-	-	-	-	-	-
2011	-	-	-	-	May (65%)	May (87%)	-	-	-
2012	May-Jun (100%)	May-Jun (67%)	Jun (89%)	Jun*(57%)	-	-	-	-	-
2013	-	-	May-Jun (89%)	Jun*(57%)	May-Jun (65%)	-	-	-	-
2014	Sep (75%)	-	-	-	-	-	Aug (35%)	-	-
2015	-	-	-	-	-	May (87%)	-	-	Aug-Sep#(17%)
2016	Aug-Sep*(100%)	Jul-Aug (67%)	Jul-Aug (89%)	Aug-Sep*(100%)	-	-	-	-	Aug-Sep#(28%)
2017	May-Jun(100%)	May-Jun (75%)	-	-	-	-	-	-	Aug# (14%)
2018	May-Jun(100%)	May-Jun (75%)	-	-	-	-	May-Jun (22%)	-	Aug# (11%)
2019	-	-	May-Jun (89%)	May-Jun (57%)	-	-	-	-	Aug-Sep#(16%)
2020	-	-	-	-	-	-	Aug-Sep	Aug-Sep	Aug-Sep
2021	-	-	Aug-Sep	Aug-Sep	Aug-Sep	Aug-Sep	-	-	-
2022	Aug-Sep	Aug-Sep	-	-	-	-	-	-	-
2023	-	-	-	-	-	-	-	-	-
2024	-	-	-	-	-	-	Aug-Sep	Aug-Sep	Aug-Sep
2025	-	-	Aug-Sep	Aug-Sep	Aug-Sep	Aug-Sep	-	-	-
2026	Aug-Sep	Aug-Sep	-	-	-	-	-	-	-
2027	-	-	-	-	-	-	-	-	-

Continue in future in the same pattern.

Table 3b

Component survey estimates to include in estimates for areas that are combinations of sub-areas. Requires updating if used in future.

	C4 = 7,8	C5 = 7WR,7E,8	C6 = 7,8,9,11	C7 = 7,8,9,11,12
1991	Yes ^a : 1990-91	Yes ^a : 1990-91	Yes ^a : 1990-91	Yes ^a : 1990-92
2003	Yes: 2002-04	Yes: 2002-04	Yes: 1999-04	Yes: 1999-04
2006	Yes ^b : 2005-07	Yes ^b : 2005-07	- (see ^c)	- (see ^c)
2013	Yes: 2012-3	Yes: 2013	Yes: 2012-14	Yes: 2012-14
2016	-	-	-	-
2017	Yes: 2016-17	Yes: 2017	Yes: 2016-18	Yes: 2016-18
2018	-	-	-	-
2019	-	-	-	-
2020	-	-	-	-
2021	Yes: 2020-21	Yes: 2021	Yes: 2020-22	Yes: 2020-22
2022	-	-	-	-
2023	-	-	-	-

Continue in future in the same pattern.

- a) The abundance estimates set for the combined sub-areas in 1990-92 assume a zero contribution from sub-area 7E as there is no available estimate for sub-area 7E to include.
- b) The abundance estimates set for combined areas C4 and C5 in 2005-07 assume a zero contribution from sub-area 7CN as there is no sub-area 7CN estimate to include.
- c) There are no 2005-2011 abundance estimate for sub-areas 9 and 12 to include in combination estimates C6 and C7; no C6 or C7 estimates are generated in this period.

Table 4

List of historical abundance estimates agreed in 2013 for use by the CLA; requires updating if they are to be used in future. Further details are given in IWC, 2014a, pp.126-9. All estimates are calculated assuming a value of 1.0 for $g(0)$ but the trials (except Trial 3) assume that $g(0) = 0.798$.

Year	SubA	Period	Est.	CV	Year	SubA	Period	Est.	CV	Year	SubA	Period	Est.	CV
1991	7CS	Aug-Sep	42*	0.603	1990	8	Aug-Sep	1,057	0.705	1990	11	Aug-Sep	2,120	0.449
2004	7CS	May	504	0.291	2002	8	Jun-Jul	63.6*	0.603	1999	11	Aug-Sep	1,456	0.565
2006	7CS	Jun-Jul	3,690	1.199	2004	8	Jun	1,093	0.576	2003	11	Aug-Sep	882	0.820
2012	7CS	May-Jun	890	0.393	2005	8	May-Jul	132	1.047	2007	11	Aug-Sep	377	0.389
1991	7CN	Aug-Sep	853	0.23	2006	8	May-Jul	309	0.677	1990	12SW	Aug-Sep	5,244	0.806
2012	7CN	Sept	398	0.507	2007	8	Jun-Jul	391	1.013	2003	12SW	Aug-Sep	3,401	0.409
1991	7WR	Aug-Sep	311	0.23	1990	9	Aug-Sep	8,264	0.396	1990	12NE	Aug-Sep	10,397	0.364
2003	7WR	May-Jun	267	0.700	2003	9	Jul-Sep	2,546	0.276	1992	12NE	Aug-Sep	11,544	0.380
2004	7WR	May-Jun	863	0.648						1999	12NE	Aug-Sep	5,088	0.377
2007	7WR	Jun-Jul	546	0.953						2003	12NE	Aug-Sep	13,067	0.287
2004	7E	May-Jun	440	0.779										
2006	7E	May-Jun	247	0.892										
2007	7E	Jun-Jul	52.6*	0.603										

The trials assume that it takes two years for the results of a sighting survey to become available to be used by a management procedure, i.e. a survey conducted in 2019 would first be used for setting the catch limit in 2021. Tables 3a and 3b list the pattern for future surveys and also show how results of surveys from different sub-areas are combined for use in variants in which *Small Areas* are comprised of more than one sub-areas. If a *Small Area* is comprised of sub-areas that are surveyed in different years, the combination abundance estimate is taken to be a summation of the estimates of abundance in the sub-areas over the years and taken to refer to the mean year (where the mean year is defined as the centre year in the set, or the later of two if this yields a half-integral year) (IWC, 1999). In cases in which the combined survey used more than one abundance estimate from the same sub-area, the abundance estimates are pooled using inverse variance weighting.

F. Parameter values and Conditioning

The biological parameters (natural mortality, age-at-maturity) and the technological parameters (selectivity) will be the same as for the previous Implementations (IWC, 1992a, p.160; IWC, 2014a, pp.133-180) (based on those for N Atlantic minke whales, IWC, 1992b, p.249)⁹ i.e.:

Table 5

The values for the biological and technological parameters that are fixed

Parameter	Value
Plus group age, x	20 yrs
Age-at-first-parturition, a_m	$m_{50} = 7$; $\sigma_m = 1.2$; first age at which a female can be mature is three,
Selectivity: Males and Females	$r_{50} = 4$; $\sigma_r = 1.2$
Maximum Sustainable Yield Level, $MSYL$	0.6 in terms of mature female component of the population

Natural mortality is age-dependent, and identical to that for the North Atlantic minke trials:

$$M_a = \begin{cases} 0.085 & \text{if } a \leq 4 \\ 0.0775 + 0.001875a & \text{if } 4 < a < 20 \\ 0.115 & \text{if } a \geq 20 \end{cases}$$

The MSYR scenarios are specified in Section G.

The ‘free’ parameters of the above model are the initial (pre-exploitation) sizes of each of the stocks, the values that determine the mixing matrices (i.e. the γ parameters), the bycatch constants (A_k). The process used to select the ‘free’ parameters is known as conditioning. The conditioning process involves first generating 100 sets of ‘target’ data as detailed in steps (a) and (b) below, and then fitting the population model to each (in the spirit of a bootstrap). The number of animals in sub-area k at the start of year t is calculated starting with guessed values of the initial population sizes and projecting the operating model forward to 2020 to obtain values of abundance etc. for comparison with the

⁹ The values are consistent with the results from JARPN. Japanese scientists advised that the above approach is appropriate given the well-known practical difficulties in using earplugs for age determination of North Pacific common minke whales. However, they also noted that technical advances mean that it may be possible to obtain age estimates in the future (IWC, 2014b, p.492).

generated data¹⁰. When performing the projections, the direct catches and known bycatches from each sub-area are set to their historical values – Appendix 1 and the bycatches are set as detailed below).

The information used in the conditioning process is as follows.

(a) Abundance estimates

The target values for the historical abundance by sub-area (except for the maximum and zero estimates – see below) are generated using the formula:

$$P_t^k = O_t^k \exp[\mu_t^k - (\sigma_t^k)^2 / 2] \quad \mu_t^k \sim N[0; (\sigma_t^k)^2] \quad (\text{F.1})$$

P_t^k is the abundance for sub-area k in year t

O_t^k is the actual survey estimate for sub-area k in year t (see Table 6); and

σ_t^k is the CV of O_t^k .

The trials are based on the two alternative values for $g(0)$ in the conditioning process: $g(0)=0.798$ (the base case value) and $g(0)=1$ (Trial 3) (IWC, 2012a, p.417; Okamura *et al.*, 2010). When $g(0)=0.798$ the values of the operating model abundances (P_t^k) are multiplied by this factor for comparison with the conditioning targets.

Minimum abundance estimates:

Table 6 includes several survey estimates that are assumed to be minima¹¹. Target values for these are similarly generated using Equation (F.1).

Maximum abundance estimates.

Bounds need to be placed on the maximum size of populations in sub-areas 5 and 6W as there is insufficient information to estimate the abundance in sub-areas 5 and 6W, given that the only estimates available for these sub-areas have very low survey coverage. Target values were generated as $P_t^k = Z_t^k / \vartheta_t^k$, where Z_t^k is the minimum estimate for the survey in the same year and period and ϑ_t^k is the proportion of the sub-area that was covered by the survey.

A maximum abundance of 500 whales in sub-area 2R in August-September 2009 was imposed in hypothesis C in the 2013 trials, to avoid undesirably high number of animals in this area. A need for such a requirement will be reviewed on inspection of the conditioning results.

Zero abundance estimates:

Table 6 includes several survey estimates of zero abundance. The target values for the historical abundance are generated using an overdispersed Poisson distribution.

(b) Proportion estimates

Estimates of the number of genetic samples assigned by stock in sub-areas 2C, 6W, 7CS, 7CN, 7WR, 10E and 11 are generated from a multinomial distribution that correspond to the observed data (see Table 7a). Some of the mixing proportions are based on data from several years so the model estimates to which these proportions are fitted during conditioning are sample size-weighted year-specific proportions.

Estimates of the proportion of recruited J-stock whales in sub-areas 6W (see Appendix 3 for how these proportions are estimated) are generated from appropriately truncated normal distributions that correspond to the observed data and are based on mtDNA and other genetic information (see Table 7b). Some of the mixing proportions are based on data from several years so year-specific proportions weighted by sample size are fitted during conditioning. A minimum standard error for the mixing proportions of 0.05 was imposed so as to prevent a few of the mixing proportions from dominating the conditioning processes – see IWC (2012c, p.106).

(c) Fixed stock proportion in sub-area 12SW

The data for sub-area 12SW are limited and so the proportion of J-stock in sub-area 12SW in June is fixed at 20% in the baseline trials. This value reflects a rough average of the J-stock mixing proportions for sub-area 11 (J-stock animals in sub-area 12SW need to pass through sub-area 11). Since the proportions for sub-area 11 are calculated from the 1984-1999 data, the 20% is taken as an average over these same years. Sensitivity trials test different levels of the sub-area 12SW proportion. In Trial 10 the proportion is 10 % (with 0% J-stock in sub-area 12NE as for the base case) and in Trial 11 the proportion is 30% (with 10% J-stock in sub-area 12NE in the same months/years; the mixing matrix is adjusted accordingly). In Trial 21 the proportion of J-stock in sub-area 12NE in May-July is fixed at 10%.

¹⁰ In order to check that the conditioning exercise has been successfully achieved, plots such as those shown in IWC (2003, pp.473-80) will be examined, together with time-trajectories of the fraction of each stock in each sub-area.

¹¹ Survey estimates based on less than 70% coverage are treated as ‘minima’ (except in sub-areas where there are no other estimates).

Table 6

Abundance data used to condition the trials**. All estimates were calculated assuming $g(0)=1$ whereas the conditioning process assumes $g(0) = 0.798$ (excepting Trial 3). See IWC, 2014a, pp. 126-9 for details of estimates used in the 2013 implementation.

Sub-area	Year	Season ^a	STD estimate ^b	CV ^c	Mode ^d	% Areal coverage	Use for Conditioning? ^e	Source
5	2001	Apr-May	1,534	0.523	NC	13	Min & Max ^f	An et al, 2010
5	2004	Apr-May	799	0.321	NC	13	Min & Max ^f	An et al, 2010
5	2008	Apr-May	680	0.372	NC	13	Min & Max ^f	An et al, 2010
5	2011	Apr-May	587	0.405	NC	13	Min & Max ^f	Park et al, 2012
6W	2000	May	549	0.419	NC	14.3	Min & Max ^f	An et al, 2010
6W	2002	May-Jun	391	0.614	NC	14.3	Min & Max ^f	An et al, 2010
6W	2003	Apr-May	485	0.343	NC	14.3	Min & Max ^f	An et al, 2010
6W	2005	Apr-May	336	0.317	NC	14.3	Min & Max ^f	An et al, 2010
6W	2006	Apr-May	459	0.516	NC	14.3	Min & Max ^f	An et al, 2010
6W	2007	Apr-May	574	0.437	NC	14.3	Min & Max ^f	An et al, 2010
6W	2009	Apr-May	884	0.286	NC	14.3	Min & Max ^f	An et al, 2010
6W	2010	Apr-May	1,014	0.387	NC	23.6	Min & Max ^f	An et al, 2011
6E	1992	Aug-Sep	893	0.67		56.8	Yes	Miyashita & Shimada, 1994
6E	2002	May-Jun	891	0.608	NC	79.1	Yes	Miyashita et al, 2009
6E	2003	May-Jun	935	0.357	NC	79.1	Yes	Miyashita et al, 2009
6E	2004	May-Jun	727	0.372	NC	79.1	Yes	Miyashita et al, 2009
10W	2006	May-Jun	2,476	0.312	IO-PS	59.9	Yes	Miyashita & Okamura 2011
10E	1992	Aug-Sep	707	0.57		30.0	Yes	Miyashita & Shimada, 1994
10E	2002	May-Jun	1,192	0.658	NC	100	Yes	Miyashita et al, 2009
10E	2003	May-Jun	591	0.566	NC	100	Yes	Miyashita et al, 2009
10E	2005	May-Jun	875	0.441	NC	64	Min	IWC, 2014a, pp.126-9
10E	2007	Jun	672	0.327	IO-PS	80.1	Yes	Miyashita et al, 2009
10E	2014	Sep	872	0.585		100	Yes	Miyashita, 2019
10E	2018	May-Jun	620	0.478		100	Yes	Hakamada et al, 2019
7CS	2004	May	504	0.291	NC	36.7	Min	IWC, 2014a, pp.126-9, 181
7CS	2006	Jul	3,690	1.199	NC	100	Yes	Hakamada & Kitakado, 2010
7CS	2012	May-Jun	537	0.346		100	Yes	Hakamada et al, 2016
7CS	2016	Aug	0	-		100	Yes	Hakamada et al, 2019
7CS	2017	May	284	0.497		100	Yes	Hakamada et al, 2019
7CS	2018	May-Jun	245	0.828		100	Yes	Hakamada et al, 2019
7CN	2012	May	542	0.601		66.7	Min	Hakamada & Kitakado, 2010
7CN	2012	Sep	599	0.525		66.7	Min	Hakamada et al, 2016
7CN	2014	Sep	244	0.454		75	Yes	Hakamada et al, 2016
7CN	2016	Aug	185	0.423		66.7	Yes	Miyashita, 2019
7CN	2017	May	179	0.377		75	Yes	Hakamada et al, 2019
7CN	2018	May	212	0.784		75	Yes	Hakamada et al, 2019
7WR	2003	May-Jun	267	0.7	NC	26.7	Min	IWC, 2014a, pp.126-9
7WR	2004	May-Jun	863	0.648	NC	88.8	Yes	Hakamada & Kitakado, 2010
7WR	2007	Jun-Jul	546	0.953		88.8	Yes	Hakamada & Kitakado, 2010
7WR	2012	Jun	378	0.79		88.8	Yes	Hakamada & Matsuoka 2016
7WR	2013	May-Jun	65	1.007		89	Yes	Hakamada et al, 2019
7WR	2016	Aug	75	1.062		89	Yes	Hakamada et al, 2019
7W: 7CS+7CN+7WR	1991	Aug-Sep	1,164	0.183			Yes	Butterworth & Miyashita, 2014
7E	2004	Jun	440	0.779	NC	57.1	Yes	Hakamada & Kitakado, 2010
7E	2006	May-Jun	247	0.892	NC	57.1	Yes	Hakamada & Kitakado, 2010
7E	2007	Jun-Jul	0	-		57.1	Yes	Hakamada & Kitakado, 2010
7E	2012	Jun	0	-		57.1	Yes	Hakamada & Matsuoka 2016
7E	2013	Jun	0	-		57.1	Yes	Hakamada et al, 2019
7E	2016	Aug	0	-		57.1	Yes	Hakamada et al, 2019
8	1990	Aug	1,057	0.706	NC	62.2	Yes	Buckland et al, 1992; Miyashita pers com 2021
8	2002	Jun-Jul	0	-	NC	65	Yes	Hakamada & Kitakado, 2010
8	2004	Jun	1,093	0.576	NC	40.5	Min	Hakamada & Kitakado, 2010
8	2005	May-Jul	132	1.047	NC	65	Yes	Hakamada & Kitakado, 2010
8	2006	May-Jul	309	0.677	NC	65	Yes	Hakamada & Kitakado, 2010
8	2007	Jun-Jul	391	1.013		65	Yes	
8	2008	Jul-Aug	0	-		65	Yes	Hakamada & Matsuoka 2016
8	2009	May-Jun	602	0.725		65	Yes	Hakamada & Matsuoka 2016
8	2011	May	121	0.966		65	Yes	Hakamada & Matsuoka 2016
8	2013	May-Jun	413	0.586		65	Yes	Hakamada et al, 2019
9	1990	Aug	3,287	0.819	NC	61.4	Yes	Buckland et al, 1992; Miyashita pers com 2021
9	2003	Jul-Sep	2,546	0.276	NC	33.2	Min	Hakamada & Kitakado, 2010
9	2008	Jul-Aug	2,458	0.664		87	Yes	Hakamada et al, 2016
9	2009	May-Jun	2,079	0.688		63	Yes	Hakamada et al, 2016
9	2011	May	115	1.025		87	Yes	Hakamada et al, 2016
9	2015	May	140	0.963		87	Yes	Hakamada et al, 2019
9N	2005	Aug-Sep	420	0.969	IO-PS	67.8	Yes	Miyashita & Okamura 2011
9N	2011	May-Jun			115	1.05	Yes	Hakamada et al, 2016

Table 6 continued

Sub-area	Year	Season ^a	STD estimate ^b	CV ^c	Mode ^d	% Areal coverage	Use for Conditioning? ^e	Source
11	1990	Aug-Sep	2,120	0.449	NC	100	Yes	Buckland et al, 1992. IWC, 2004, p.124
11	1999	Aug-Sep	1,456	0.565	IO	100	Yes	IWC, 2004, p.124
11	2003	Aug-Sep	882	0.826	IO-AC	33.9	Min	Miyashita & Okamura, 2011
11	2007	Aug-Sep	377	0.389	IO-PS	20.2	Min	Miyashita & Okamura, 2011
11	2014	Aug	306	0.679		35	Min	Miyashita, 2019
11	2018	May	235	0.481		21.7	Min	Hakamada et al, 2019
12SW	1990	Aug-Sep	4,774	0.508	NC	100	Yes	Buckland et al, 1992. cv recalculated (Miyashita pers. comm 2021).
12SW	2003	Aug-Sep	3,401	0.409	IO-AC	100	Yes	Miyashita & Okamura, 2011
12NE	1990	Aug-Sep	11,805	0.377	NC	100	Yes	Buckland et al, 1992. Recalculated Miyashita pers. comm Nov 2021
12NE	1992	Aug-Sep	11,051	0.705	NC	[100]	Yes	Miyashita & Shimada, 1994; Recalculated Miyashita pers. comm Nov 2021
12NE	1999	Aug-Sep	5,088	0.377	NC	63.8	Min	IWC, 2014a, pp.126-9
12NE	2003	Aug-Sep	13,067	0.287	IO-AC	41	Min	Miyashita & Okamura, 2011

** The above table lists estimates used in conditioning, including corrections received from Japan. The Secretariat maintains a full list of estimates including details of other estimates and the reason they were not included in the above table.

^a Season: if a survey took place in less than 20% of a month, that month was not used as part of the survey-time-period in the likelihood calculation.

^b Standard (STD) estimate based on 'Top and Upper bridge' assuming g(0)=1, but subsequently corrected by estimate of g(0) for the combined platform 'Top and Upper bridge'.

^c CV does not consider any process errors.

^d Mode: NC=Normal-closing, IO-PS=Passing with IO mode, IO-AC=Abeam-closing with IO mode. (STD estimates by different modes, NC, IO-AC, IO-NC, are considered comparable.)

^e Survey estimates based on less than 70% coverage are treated as 'minima' (except in sub-areas where there are no other estimates).

^f Maximum values are calculated as the best estimate / coverage.

(d) Fixed stock proportion in sub-area 9 and 9N

The data for sub-area 9 are also limited. For Trials 2 and 23, which assume a C-stock that mixes with the O-stock in sub-area 9 and 9N, the proportion of O-stock is assumed to be 0.5 during August and September in 1995. This is based on the ratio assumed in 9W in 2003. For hypothesis E, Trial 2 the same proportion is also assumed in 12NE in August and September 1995 (but not in Trial 23).

Table 7a

The number of sampled whales that were assigned to each stock using the genetic assignment data based on STRUCTURE (Hypothesis A & B) and Geneland (Hypothesis E) using a 90% probability of assignment, except for Trial 5 where a 70% probability of assignment is used. In sub-areas 7CS and 7CN the baseline and Trial 5 proportion of whales assigned to each stock is weighted by 5/60 of the bycatch proportion and 55/60 of the special permit proportion. The number assigned by stock is then taken as this proportion multiplied by the total number of assigned animals. In Trial 6 the proportion of whales assigned to each stock is weighted by 2/60 of the bycatch proportion and 58/60 of the special permit proportion, while in Trial 7 10/60 of the bycatch proportion and 50/60 of the special permit proportion are used. These data are used to condition the trials.

Hypothesis	Trial	Area	Years	Months	Sex	Total Sample	Bycatch Samples		Special Permit Samples		Weighted Total	
							J-Stock	O-Stock	J-Stock	O-Stock	J-Stock	O-Stock
A & B	Baseline	2C	2002-16	Jan-Apr	M+F	155	127	28			127	28
A & B	Baseline	2C	2001-16	May-Sep	M+F	56	46	10			46	10
A & B	Baseline	2C	2001-16	Oct-Dec	M+F	134	122	12			122	12
A & B	Baseline	7CS	2002-16	Jan-Apr	M+F	263	43	34	48	138	74	189
A & B	Baseline	7CS	2001-16	May	M+F	391	16	31	89	255	104	287
A & B	Baseline	7CS	1999-2016	Jun-Dec	M+F	199	86	34	4	75	21	178
A & B	Baseline	7CN	2002-16	Jan-May	M+F	100	27	29	6	38	17	83
A & B	Baseline	7CN	1999-2016	Jun	M+F	133	11	15	6	101	12	121
A & B	Baseline	7CN	1996-2016	Jul-Sep	M+F	610	16	13	103	478	127	483
A & B	Baseline	7CN	2001-16	Oct-Dec	M+F	270	35	2	66	167	91	179
A & B	Baseline	10E	2001-16	Jun-Dec	M+F	15	14	1			14	1
A & B	Baseline	11	1996-2012	May-Dec	M	57					28	29
A & B	Baseline	11	1996-2015	May-Dec	F	58					28	30
Hypothesis	Trial	Area	Years	Months	Sex	Total Sample	J-Stk	P-Stk	O-Stk	J-Stk	P-Stk	O-Stk
E	Baseline	2C	2002-16	Jan-Apr	M+F	138	107	31			107	31
E	Baseline	2C	2001-16	May-Sep	M+F	49	32	17			32	17
E	Baseline	2C	2001-16	Oct-Dec	M+F	122	105	17			105	17
E	Baseline	7CS	2002-16	Jan-Apr	M+F	262	-	73	1	-	188	-
E	Baseline	7CS	2001-16	May	M+F	378	-	49	2	-	303	24
E	Baseline	7CS	1999-2016	Jun-Dec	M+F	197	-	118	1	-	5	73
E	Baseline	7CN	1999-2016	Jan-Jun	M+F	220	12	69	-	2	28	109
E	Baseline	7CN	1996-2016	Jul-Dec	M+F	881	13	59	-	10	574	225
E	Baseline	11	1996-2012	May-Dec	M	59					13	45
E	Baseline	11	1996-2015	May-Dec	F	63					18	41

Hypothesis	Trial	Area	Years	Months	Sex	Total Sample	Bycatch Samples		Special Permit Samples		Weighted Total	
							J-Stock	O-Stock	J-Stock	O-Stock	J-Stock	O-Stock
A & B	5	2C	2002-16	Jan-Apr	M+F	170	138	32			138	32
A & B	5	2C	2001-16	May-Sep	M+F	57	47	10			47	10
A & B	5	2C	2001-16	Oct-Dec	M+F	141	129	12			129	12
A & B	5	7CS	2002-16	Jan-Apr	M+F	291	48	41	51	151	80	211
A & B	5	7CS	2001-16	May	M+F	431	17	37	100	277	116	315
A & B	5	7CS	1999-2016	Jun-Dec	M+F	212	92	36	4	80	22	190
A & B	5	7CN	2002-16	Jan-May	M+F	105	28	30	7	40	19	86
A & B	5	7CN	1999-2016	Jun	M+F	139	12	16	8	103	14	125
A & B	5	7CN	1996-2016	Jul-Sep	M+F	660	20	14	109	517	138	522
A & B	5	7CN	2001-16	Oct-Dec	M+F	283	38	2	67	176	94	189
A & B	5	7WR+7E	1996-2006	May	M+F	87					3	84
A & B	5	7WR+7E	1996-2012	Jun-Aug	M+F	49					0	49
A & B	5	8	1998-2012	May-Jun	M+F	139					1	138
A & B	5	8	1996-2009	Jul-Sep	M+F	106					1	105
A & B	5	9	1995-2011	May-Jun	M+F	125					1	124
A & B	5	9	1994-2010	Jul	M+F	190					4	186
A & B	5	9	1994-2013	Aug-Sep	M+F	212					0	212
A & B	5	10E	2001-16	Jun-Dec	M+F	16					15	1
A & B	5	11	1996-2012	May-Dec	M	64					30	34
A & B	5	11	1996-2015	May-Dec	F	63					30	33
A & B	6	7CS	2002-16	Jan-Apr	M+F	263	43	34	48	138	71	192
A & B	6	7CS	2001-16	May	M+F	391	16	31	89	255	102	289
A & B	6	7CS	1999-2016	Jun-Dec	M+F	199	86	34	4	75	14	185
A & B	6	7CN	2002-16	Jan-May	M+F	100	27	29	6	38	15	85
A & B	6	7CN	1999-2016	Jun	M+F	133	11	15	6	101	9	124
A & B	6	7CN	1996-2016	Jul-Sep	M+F	610	16	13	103	478	116	494
A & B	6	7CN	2001-16	Oct-Dec	M+F	270	35	2	66	167	82	188
A & B	7	7CS	2002-16	Jan-Apr	M+F	263	43	34	48	138	81	182
A & B	7	7CS	2001-16	May	M+F	391	16	31	89	255	106	285
A & B	7	7CS	1999-2016	Jun-Dec	M+F	199	86	34	4	75	32	167
A & B	7	7CN	2002-16	Jan-May	M+F	100	27	29	6	38	19	81
A & B	7	7CN	1999-2016	Jun	M+F	133	11	15	6	101	16	117
A & B	7	7CN	1996-2016	Jul-Sep	M+F	610	16	13	103	478	146	462
A & B	7	7CN	2001-16	Oct-Dec	M+F	270	35	2	66	167	106	144
Hypothesis	Trial	Area	Years	Months	Sex	Total Sample	J-Stk	P-Stk	O-Stk	J-Stk	P-Stk	O-Stk
E	5	2C	2002-16	Jan-Apr	M+F	150					116	33
E	5	2C	2001-16	May-Sep	M+F	54					36	18
E	5	2C	2001-16	Oct-Dec	M+F	125					108	17
E	5	7CS	2002-16	Jan-Apr	M+F	282	2	79	3	2	196	0
E	5	7CS	2001-16	May	M+F	411	0	51	2	1	326	31
E	5	7CS	1999-2016	Jun-Dec	M+F	211	1	127	2	0	8	73
E	5	7CN	1999-2016	Jan-Jun	M+F	237	13	70	-	2	30	122
E	5	7CN	1996-2016	Jul-Dec	M+F	915	15	59	-	11	582	248
E	5	11	1996-2012	May-Dec	M	63					14	48
E	5	11	1996-2015	May-Dec	F	64					18	42
E	6	7CS	2002-16	Jan-Apr	M+F	262	-	73	1	-	188	-
E	6	7CS	2001-16	May	M+F	378	-	49	2	-	303	24
E	6	7CS	1999-2016	Jun-Dec	M+F	197	-	118	1	-	5	73
E	6	7CN	1999-2016	Jan-Jun	M+F	220	12	69	-	2	28	109
E	6	7CN	1996-2016	Jul-Dec	M+F	881	13	59	-	10	574	225
E	7	7CS	2002-16	Jan-Apr	M+F	262	-	73	1	-	188	-
E	7	7CS	2001-16	May	M+F	378	-	49	2	-	303	24
E	7	7CS	1999-2016	Jun-Dec	M+F	197	-	118	1	-	5	73
E	7	7CN	1999-2016	Jan-Jun	M+F	220	12	69	-	2	28	109
E	7	7CN	1996-2016	Jul-Dec	M+F	881	13	59	-	10	574	225

Table 7b

Estimates of the proportion of recruited 'J'-whales used to condition the trials based on mtDNA and Allele samples.

Hypothesis	Area	Years	Months	Sex	Ratio	CV ¹²	Data Type	Stock
B and E	6W	1999-2007	Jan-Mar	M+F	0.584	0.131	mtDNA	J:Total
B and E	6W	1999-2007	Jan-Mar	M+F	0.672	0.05	Allelle	J:Total
B and E	6W	1999-2007	Apr-Jun	M+F	0.496	0.126	mtDNA	J:Total
B and E	6W	1999-2007	Apr-Jun	M+F	0.812	0.05	Allelle	J:Total
B and E	6W	1999-2007	Jul-Aug	M+F	1.000	0.05	mtDNA	J:Total
B and E	6W	1999-2007	Jul-Aug	M+F	0.749	0.077	Allelle	J:Total
B and E	6W	1999-2007	Sep-Dec	M+F	0.593	0.123	mtDNA	J:Total
B and E	6W	1999-2007	Sep-Dec	M+F	0.761	0.05	Allelle	J:Total
B and E	6W	1999-2007	Sep-Dec	M+F			Bycatch samples	

¹² In cases when the sample size used to generate the proportion estimates is small and the se's are small (which will overweight such results), the standard error is set to 0.05.

(f) Calculation of likelihood

The objective function consists of three components: Objective Function = $-(L_1+L_2+L_3)$. Equations F.4-6 list the negative of the logarithm of the objective function for each of the three components:

Abundance estimates

$$L_{1a} = 0.5 \sum_n \frac{1}{(\sigma_t^k)^2} \left(\ln \left(P_n^k / \hat{P}_n^k \right) \right)^2 \quad (\text{F.4a})$$

where \hat{P}_n^k is the model estimate of the abundance in the same year, period and sub-area as the n th estimate of abundance P_n^k .

Minimum abundance estimates

$$L_{1b} = \sum_n \left\{ \ln \sigma_t^k + \frac{1}{2(\sigma_t^k)^2} \ln \left(P_n^k / \hat{P}_n^k \right)^2 \right\} \left\{ \frac{\exp(\Delta(P_n^k - \hat{P}_n^k))}{1 + \exp(\Delta(P_n^k - \hat{P}_n^k))} \right\} + \ln \sigma_t^k \left\{ \frac{1}{1 + \exp(\Delta(P_n^k - \hat{P}_n^k))} \right\} \quad (\text{F.4b})$$

where Δ is a “large” number (here 30).

Maximum abundance estimates

$$L_{1c} = \sum_n \left\{ \ln \sigma_t^k + \frac{1}{2(\sigma_t^k)^2} \ln \left(P_n^k / \hat{P}_n^k \right)^2 \right\} \left\{ \frac{1}{1 + \exp(\Delta(P_n^k - \hat{P}_n^k))} \right\} + \ln \sigma_t^k \left\{ \frac{\exp(\Delta(P_n^k - \hat{P}_n^k))}{1 + \exp(\Delta(P_n^k - \hat{P}_n^k))} \right\} \quad (\text{F.4c})$$

Zero abundance estimates

$$L_{1d} = - \sum_n [n_n^k \ln(\beta_n^k \hat{P}_n^k) - \beta_n^k \hat{P}_n^k] / \hat{\alpha}_n^k \quad (\text{F.4d})$$

where n_n^k is the number of animals seen during the n th survey in sub-area k , β_n^k is the realised track length for the n th survey in sub-area k multiplied by the average effective search half width, and divided by the sub-area size (Table 8), \hat{P}_n^k is the model-estimate corresponding to the n th survey in sub-area k and $\hat{\alpha}^k$ is the adjusted coefficient of variation of the survey estimate P_n^k , $\hat{\alpha}^k = \frac{\sum_m (n_m^k)^2 CV^2(P_m^k)}{\sum_m n_m^k}$, constrained to $\hat{\alpha}^k \geq 1$, where m denotes the number of (non-minima) survey estimates within sub-area k for which the number of animals seen and the CV of the survey estimate are available. See Appendix 4 for the derivation of this equation.

Table 8

The realised track length, average effective search half width and sub-area size corresponding to the zero abundance estimates. The effective search half width is taken to be the average from other surveys (excluding those considered minimum estimates) in the same sub-area used in conditioning, for which effective search half width is available.

Year	Sub-Area	Realised track length	Average effective search half width [No. of surveys used]	Sub-area size	$\hat{\alpha}^k$
2016	7CS	754	0.3955 [4]	26826	22.83
2007	7E	360	0.4225 [2]	84427	1.73
2012	7E	302	0.4225 [2]	84427	1.73
2013	7E	599	0.4225 [2]	84427	1.73
2016	7E	472	0.4225 [2]	84427	1.73
2008	7	887	0.374 [1]	217678	1.00 ¹³
2002	8	1184	0.5283 [7]	250291	1.50
2008	8	1194	0.5283 [7]	250445	1.50

Stock proportions

For sub-areas 2C, 7CN, 7CS, 10E and 11:

$$L_2 = - \sum_j \sum_n N_{j,n}^k \ln(\hat{p}_{j,n}^k / p_{j,n}^{obs,k}) \quad (\text{F.5a})$$

where $\hat{p}_{j,n}^k$ is the model estimate of the proportion of j -stock whales in the same year, period, sub-area and gender as the n th set of data and $p_{j,n}^{obs,k}$ is the corresponding observed value, with $N_{j,n}^k$ denoting the observed number of samples of j -stock whales in the n th set of data.

¹³ Due to constraint of $\hat{\alpha}^k \geq 1$

For sub-area 6W in Hypotheses B and E only:

$$L_2 = 0.5 \sum_n \frac{1}{(\sigma_n^k)^2} (p_n^k - \hat{p}_n^k)^2 \quad (\text{F.5b})$$

where \hat{p}_n^k is the model estimate of the proportion of whales in the same year, period and sub-area as the n th proportion estimate p_n^k .

Bycatch estimates

$$L_3 = 0.5 \sum_n (B_n^k - \hat{B}_n^k)^2 / 10 \quad (\text{F.6})$$

where \hat{B}_n^k is the model estimate of the total bycatch in sub-area k over the years being fitted and B_n^k is the observed bycatch in the same area and period.

G. Trials

The factors to be considered based on the previous trials are listed in Table 9 and the set of trials in Table 10. The sensitivity trials are variants of the base-case trials A01-1 etc. (see section A).

H. Management options

Future direct catch options will be specified later.

I. Output statistics

Population-size and continuing catch statistics are produced for each stock, and catch-related statistics for each sub-area. Catch-related statistics are produced both for the total catches (commercial and incidental) and for the commercial catches alone.

- (1) Total catch (TC) distribution: (a) median; (b) 5th value; (c) 95th value.
- (2) Initial mature female population size (P_{1930}) distribution: (a) median; (b) 5th value; (c) 95th value.
- (3) Final mature female population size (P_{2120}) distribution: (a) median; (b) 5th value; (c) 95th value.
- (4) Lowest mature female population size over 100 years (P_{low}) distribution: (a) median; (b) 5th value; (c) 95th value.
- (5) Average catch over the last 10 years of the 100-year management period: (a) median; (b) 5th value; (c) 95th value.
- (6) Catch by sub-area, stock and catch-type (incidental or commercial): (a) median; (b) 5th value; (c) 95th value.
- (7) The median percentage of mature J-stock females being in sub-area 12 in June-August 1973-75.
- (8) The median annual rate of decline in the number of whales assumed recruited to the Korean fishery over the period 1973-1986.
- (9) The median 1+ population size for animals in sub-areas 6 and 10 in August-September in 1992 and in 2000 (corresponding to Sea of Japan surveys).
- (10) Proportion Mature: compare the numbers of mature animals by sub-area and time period with the (approximate) proportion mature in the available observation data.
- (11) The mean proportion of J whales in the total (scientific, commercial and incidental) catch taken by Japan from 1993-98 is output in trials, for comparison with results obtained from market samples.

Table 9
Proposed factors to be considered in the Trials.

Factor
Stock structure hypothesis
Stock structure hypotheses A, B and E
MSYR
1% ₁₊ ; 4% _{mat}
g(0)
0.798; 1.00 (Trial 3)
Other stock structure issues
With a C-stock, i.e. from a putative 'Central' North Pacific population (Trial 2)
Alternative basis for mixing rates (Trial 5)
10% J-stock in sub-area 12SW in June (Trial 10)
30% J-stock in sub-area 12SW in June (Trial 11)
No C-stock (i.e. from a putative 'Central' North Pacific population) in sub-area 12NE (Trial 23)
10% J-stock in sub-area 12NE in May-July (Trial 21)
Catches and bycatches
High direct catches (Baseline total = 39,299; high total = 40,879) + alternative Korean & Japanese bycatch level (Trial 4)
Different allocation of the Korean catches between sub-areas 5 and 6W. (Trials 8 and 9)
Chinese incidental catch = 0 (Trial 12) (Baseline value = 2* Korean bycatch in sub-area 5)
Number of bycaught animals is proportional to square root of abundance (Trial 17)
Mixing and dispersion
Mixing proportion in sub-areas 7CS and 7CN calculated using alternative weighting for bycatch: 2/60 weight (Trial 6) and 10/60 weight (Trial 7)
A substantially larger fraction of whales aged 1-4 from O-stock are found in sub-areas 2R, 3 and 4 year round (Trial 18)
Set the proportion of O-stock animals of ages 1-4 in sub-areas 9 and 9N to zero (Trial 19)
Time-varying mixing matrix for the bycatch (Trial 22) (requires specification)
Abundance estimates
The 2013 implementation considered Alternative abundance estimates for sub-areas 6E and 10E.
The number of 1+ whales in 2009 in sub-area 2C in any month < 200 (Trial 20)

Table 10
The list of trials (MSYR 1% is defined in terms of the total (1+) component and 4% on the mature female component of the population).

Stock hypothesis	Trial no.	MSYR	Mix matrix:	Description
A	A01-1 & A01-4	1% / 4%	Baseline	Baseline A: 2 stocks (J- and O-); g(0) = 0.798; including Chinese bycatch
B	B01-1 & B01-4	1% / 4%	Baseline	Baseline B: 3 stocks (J-, O-, and Y-); g(0) = 0.798; including Chinese bycatch
E	E01-1 & E01-4	1% / 4%	Baseline	Baseline E: 5 stocks (J-, P-, O-, and Y-); g(0) = 0.798; including Chinese bycatch
AE	A02-1 etc	1% / 4%	Trial 2	With a C- ('Central' North Pacific) stock
ABE	A03-1 etc	1% / 4%	Baseline	Assume g(0) = 1
ABE	A04-1 etc	1% / 4%	Baseline	High direct catches + alternative Korean & Japanese bycatch levels
ABE	A05-1 etc	1% / 4%	Trial 5	Alternative (70% probability) thresholds for assignment of stock proportions
ABE	A06-1 etc	1% / 4%	Baseline	No. of genetic samples assigned to stock in sub-areas 7CS and 7CN calculated using 2/60 weight for bycatch
ABE	A07-1 etc	1% / 4%	Baseline	No. of genetic samples assigned to stock in sub-areas 7CS and 7CN calculated using 10/60 weight for bycatch
ABE	A08-1 etc	1% / 4%	Baseline	More Korean catches in sub-area 5 (and fewer in sub-area 6W). Rationale: the baseline uses the best split. Trials 8 and 9 test alternatives in both directions.
ABE	A09-1 etc	1% / 4%	Baseline	More Korean catches in sub-area 6W (and fewer in 5)
ABE	A10-1 etc	1% / 4%	Baseline	10% J -stock in sub-area 12SW in June (base case value = 20%). See section F(c).
ABE	A11-1 etc	1% / 4%	Trial 11	30% J -stock in sub-area 12SW in June (base case value = 20%) with 10% J -stock in 12NE in May-June. See section F(c).
ABE	A12-1 etc	1% / 4%	Baseline	Chinese incidental catch = 0 (the base case value = twice that of Korea in sub-area 5)
ABE	A13-1 etc	1% / 4%	Baseline	Alternative abundance estimates in sub-area 6E (see table 6)
ABE	A14-1 etc	1% / 4%	Baseline	Additional abundance estimate in sub-area 10E in 2007 (see table 6)
AE	A17-1 etc	1% / 4%	Baseline	The number of bycaught animals is proportional to the square-root of abundance rather than to abundance (in order to examine the impact of possible saturation effects)
ABE	A18-1 etc	1% / 4%	Trial 18	A substantially larger fraction of whales ages 1-4 from O-stock are found in sub-areas 2R, 3 and 4 year-round (so the proportion of 1-4 whales in sub-area 9 is closer to expectations given the length-frequencies of catches from sub-area 9). The mixing matrices are adjusted such that the numbers of age 1-4 of O-stock animals in sub-areas 9 and 9N are no more than half the base case numbers; juveniles are allowed into sub-areas 2R, 3 and 4 in the corresponding months.
ABE	A19-1 etc	1% / 4%	Trial 19	Set the proportion of O animals of ages 1-4 in sub-areas 9 and 9N to zero and allow the abundance in sub-areas 7CS and 7CN to exceed the abundance estimates for these sub-areas. Projections for these sub-areas will need to account for the implied survey bias
ABE	A20-1 etc	1% / 4%	Trial 20	The number of 1+ whales in 2009 in sub-area 2C in any month < 200 (if large numbers of whales were found in 2C, the historical catch would be expected to be much greater).
ABE	A21-1 etc	1% / 4%	Trial 21	10% J -stock in sub-area 12NE in May-July. See section F(c).
ABE	A22-1 etc	1% / 4%	Trial 22	Time-varying mixing matrix for the bycatch
E	E23-1 & 4	1% / 4%	Trial 23	With a putative C ('Central North' Pacific) stock, but no C animals in sub-area 12NE

REFERENCES

- Allison, C. 2011. Direct catch data for western North Pacific minke whale simulation trials. Paper SC/D11/NPM3 presented to the First Intersessional Workshop for the *Implementation Review* of western North Pacific common minke whales, 12-16 December 2011, Tokyo, Japan (unpublished). [Paper available from the Office of this Journal].
- An, Y.R., Choi, C.D., Moon, D.Y. and Park, K.H. 2010. Summary of the information on Korean dedicated sighting surveys for abundance estimates of common minke whales. Paper SC/D10/NPM15 presented to the First Intersessional Workshop for Western North Pacific Common Minke Whales, 14-17 December 2010, Pusan, Republic of Korea (unpublished). 7pp. [Paper available from the Office of this Journal].
- An, Y.-R., Park, K.-Y., Choi, S.-G., Kim, H.-W. And Moon, D.-Y. 2011. Abundance estimation of western North Pacific minke whales using the Korean sighting survey in 2010. 4pp.
- Baker, C.S., Cooke, J.G., Lavery, S., Dalebout, M.L., Ma, Y.U., Funahashi, N., Carraher, C. and Brownell, J., R.L. 2007. Estimating the number of whales entering trade using DNA profiling and capture-recapture analysis of market produce. *Mol. Ecol.* 16(13): 2617-26.
- Butterworth D. and Miyashita T. 2014. Derivation of revised estimate for subarea 7 in 1991 and zero abundance estimates. Annex F. *J. Cetacean Res. Manage. (Suppl.)* 15:502-3.
- Hakamada, T. 2010. The number of set nets in the coast of Japan by sub-areas and years during 1979-2006. Paper SC/D10/NPM13 presented to the First Intersessional Workshop for Western North Pacific Common Minke Whales, 14-17 December 2010, Pusan, Republic of Korea (unpublished). 1pp. [Paper available from the Office of this Journal].
- Hakamada, T. and Kitakado, T. 2010. Abundance estimation for western North Pacific common minke whales using sighting information from JARPN and JARPN II. Paper SC/D10/NPM12rev 25pp. [Paper available from the Office of this Journal].
- Hakamada, T. and Matsuoka, K. 2016. The number of western North Pacific common minke, Bryde's and sei whales distributed in JARPN II Offshore survey area. Paper SC/F16/JR12. 13pp. [Paper available from the Office of this Journal].
- Hakamada, T., Matsuoka, K., Toshiya, K. and Miyashita, T. 2016. The number of the western North Pacific common minke whales (*Balaenoptera acutorostrata*) distributed in JARPN II coastal survey areas. Paper SC/F16/JR11. 7pp. [Paper available from the Office of this Journal].
- Hakamada, T., Katsumata, T., Takahashi, M. and Matsuoka, K. 2019. Common minke whale abundance estimates based on dedicated sightings surveys during 2013-2018. Paper SC68a/ASI/14rev (unpublished). [Paper available from the Office of this Journal].
- International Whaling Commission. 1991. Report of the Sub-Committee on Management Procedures, Appendix 4. Report of the *ad-hoc* trials subgroup. *Rep. int. Whal. Commn* 41:108-12.
- International Whaling Commission. 1992a. Report of the Scientific Committee, Annex F. Report of the Sub-Committee on North Pacific Minke Whales. *Rep. int. Whal. Commn* 42:156-77.
- International Whaling Commission. 1992b. Report of the Scientific Committee, Annex K. Report of the Working Group on North Atlantic Minke Trials. *Rep. int. Whal. Commn* 42:246-51.
- International Whaling Commission. 1994. Report of the Scientific Committee, Annex D. Report of the Sub-Committee on Management Procedures, Appendix 2. Minimum Standards Trials. *Rep. int. Whal. Commn* 44:85-88.
- International Whaling Commission. 1999. Report of the Scientific Committee. Annex N. The Revised Management Procedure (RMP) for Baleen Whales. *J. Cetacean Res. Manage. (Suppl.)* 1:251-58.
- International Whaling Commission. 2004. Report of the Scientific Committee. Annex D. Report of the Sub-Committee on the Revised Management Procedure. *J. Cetacean Res. Manage. (Suppl.)* 6: 75-184.
- International Whaling Commission. 2012a. Report of the first RMP intersessional workshop for western North Pacific common minke whales. *J. Cetacean Res. Manage. (Suppl.)* 13:411-60.
- International Whaling Commission. 2012b. Report of the first RMP intersessional workshop for western North Pacific common minke whales. Annex G. NPM mixing matrices. *J. Cetacean Res. Manage. (Suppl.)* 13:445-48.
- International Whaling Commission. 2012c. Report of the Scientific Committee. Annex D1. Report of the Working Group on the *Implementation Review* for western North Pacific common minke whales. *J. Cetacean Res. Manage. (Suppl.)* 13:102-29.
- International Whaling Commission. 2014a. Report of the Scientific Committee. Annex D1. Report of the Working Group on the *Implementation Review* for western North Pacific common minke whales. *J. Cetacean Res. Manage. (Suppl.)* 15: 112-88.
- International Whaling Commission. 2014b. Report of the 'Second' Intersessional Workshop on the Implementation Review for western North Pacific Common Minke Whales, 19-23 March 2013, La Jolla, California, USA. *J. Cetacean Res. Manage. (Suppl.)* 15:489-506.
- International Whaling Commission. 2020. Report of the 'First' Intersessional Workshop on the Implementation Review for western North Pacific Common Minke Whales, 25 February-1 March 2019, Tokyo, Japan. *J. Cetacean Res. Manage. (Suppl.)* 21:373-445.
- Miyashita, T. 2010. Summary of the information on Japanese dedicated sighting surveys for abundance estimation. Paper SC/D10/NPM11 presented to the First Intersessional Workshop for Western North Pacific Common Minke Whales, 14-17 December 2010, Pusan, Republic of Korea (unpublished). 6pp. [Paper available from the Office of this Journal].
- Miyashita, T. and Okamura, H. 2011. Abundance estimates of common minke whales using the Japanese dedicated sighting survey data for RMP Implementation and CLA - Sea of Japan and Sea of Okhotsk. Paper SC/63/RMP11 presented to the IWC Scientific Committee, June 2011, Tromsø, Norway (unpublished). 34pp.
- Miyashita, T., Okamura, H. and Kitakado, T. 2009. Abundance of J-stock common minke whales in the Sea of Japan using the Japanese sighting data with $g(0)=1$. Paper SC/61/NPM7 presented to the IWC Scientific Committee, June 2009, Madeira, Portugal (unpublished). 10pp.
- Miyashita, T. and Shimada, H. 1994. Minke whale abundance in the Okhotsk Sea, the Sea of Japan and off the Pacific coast of Northern Japan estimated from sighting data. Paper SC/46/NP6 presented to the IWC Scientific Committee, May 1994 (unpublished). 9pp. [Paper available from the Office of this Journal].
- Miyashita, 2019. Abundance estimate of common minke whales in sub-areas 11, 10E and 7CN in 2014. Paper SC/68a/ASI 15. [Paper available from the Office of this Journal].
- Okamura, H., Miyashita, T. and Kitakado, T. 2010. $g(0)$ estimates for western North Pacific common minke whales. Paper SC/62/NPM9 presented to the IWC Scientific Committee, June 2010, Agadir, Morocco (unpublished). 7pp. [Paper available from the Office of this Journal].
- Park, K.J., An, Y.R., Kim, H.W., Kim, D.N., Sohn, H.S. And An, D.H. 2012. Abundance estimation of common minke whales in the Yellow Sea using the Korean sighting data in 2011. 4pp. Paper SC/64/NPM7. [Paper available from the Office of this Journal].
- Punt, A.E. 1999. Report of the Scientific Committee. Annex R. A full description of the standard BAILEN II model and some variants thereof. *J. Cetacean Res. Manage. (Suppl.)* 1: 267-76.
- Tobayama, T., Yanagisawa, F. and Kasuya, T. 1992. Incidental take of minke whales in Japanese trap nets. *Rep. int. Whal. Commn* 42: 433-36.

Appendix 1

The Historical Catch Series

C. Allison

Direct catches

The baseline trials use the 'best' estimates of the historical direct catch, which are summarised in Tables 1 and 2. Details of the sources and construction of the catch series are given in Allison (2011). The data are taken from the IWC individual catch database (Allison, 2020) where available; where these data are not available the catch series has been compiled to match all known sources of information.

An alternative 'high' catch series is used in Trial 4. Table 3 lists the 'high' catch numbers for the years and sub-areas where they differ from the 'best' catch series. The catches are identical to the 'best' series for all other areas and years.

The Japanese coastal catch from 1930-1 and 1936-45 (in sub-areas 7CS, 7CN and 11) is estimated (Ohsumi 1982) and the values are doubled in the 'high' catch series. The catch series off Korea assumes a linear increase from 60 whales in 1946 to 249 in 1957 in the 'best' series whereas the 'high' series assumes an annual catch of 249 minke whales over this period.

The split between sub-areas 5 and 6W is unknown for most of the catches taken off Korea. The 'best' catch series includes 19,349 minke whales taken off Korea, of which 3,902 are recorded in the Yellow Sea and 4,199 in the Sea of Japan (East Sea) and Southern waters. The remaining 11,248 of unknown area are allocated between sub-areas 5 and 6W in the ratio of the catches known by area from 1940-79¹⁴ (2,028:2,517). Where catches are known by month from 1958-86, (Park 1995) but not area, they are allocated to sub-area using the average known ratio in the given month. Trials 8 and 9 test the sensitivity to this assumption. In Trial 8 the number of whales allocated to sub-area 5 is reduced by 20% and reallocated to sub-area 6W. In Trial 9, 20% fewer animals are allocated to sub-area 6W and are reallocated to sub-area 5. The resulting catch series is given in Table 4.

Table 1.

Summary of the final western North Pacific Minke Whale Direct Catch Series (1930-2020) by sub-area, sex and month. Catches that cannot be taken because no whales are modelled the area/month are highlighted.

Area	Males							Females							Total	M	F		
	J-M	Apr	May	Jun	Jul	Aug	Sep	O-D	J-M	Apr	May	Jun	Jul	Aug	Sep	O-D			
1E	17	0	0	0	1	0	0	0	11	0	0	0	0	0	0	0	29	18	11
2C	'3	2	2	3	2	0	1	0	2	2	0	0	1	0	0	0	18	13	5
2R	1	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	4	2	2
5	981	1,280	906	671	568	322	102	174	1,128	1,457	1,244	757	570	300	121	185	10,766	5,004	5,762
6W	181	383	1,325	1,167	392	202	557	1,063	178	364	1,300	1,136	376	189	545	1,009	10,367	5,270	5,097
6E	181	223	135	13	21	0	8	2	95	144	95	16	3	0	6	1	943	583	360
7CS	210	1,011	1,826	768	129	8	1	0	164	1,134	1,371	464	27	1	0	0	7,114	3,953	3,161
7CN	0	0	77	241	387	426	940	199	0	20	89	101	163	122	312	113	3,190	2,270	920
7W	0	1	49	33	3	1	10	0	0	0	9	3	3	0	0	0	112	97	15
7E	0	0	37	21	3	0	13	1	0	0	7	2	0	0	9	0	93	75	18
8	0	0	39	101	99	21	11	6	0	0	8	10	17	4	5	6	327	277	50
9	0	0	32	82	183	218	17	0	0	0	9	11	16	29	3	0	600	532	68
9N	0	0	1	2	5	8	0	1	0	0	0	6	0	11	0	0	34	17	17
10W	0	0	6	12	1	0	2	0	0	2	0	9	0	0	0	0	32	21	11
10E	2	25	42	119	83	26	5	3	0	1	28	60	26	9	7	0	436	305	131
11	0	62	248	503	560	230	143	29	2	465	872	909	607	273	113	25	5,041	1,775	3,266
12SW	0	0	0	1	11	9	1	0	0	0	1	5	16	27	5	0	76	22	54
12NE	0	0	0	0	36	9	10	0	0	0	0	3	33	14	6	0	111	55	56
13	0	0	0	0	0	2	0	0	0	0	0	1	3	0	0	0	6	2	4
Total	1,576	2,988	4,725	3,737	2,484	1,482	1,821	1,478	1,581	3,589	5,033	3,492	1,859	982	1,133	1,339	39,299	20,291	19,008

¹⁴The period 1940-79 is used in view of a comment by Gong (1982) that, in 1980, Government policy led to a shift to the western sector in order to direct the minke whale fishery away from areas where the (protected) fin whale might also be caught.

Table 2
Summary of the 'Best' Direct Catch Series for western North Pacific Minke Whales by Year, sub-area and sex.

Males:

	1E	2C	2R	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13	Total	
1930	0	0	0	0	0	0	7	0	0	0	0	0	0	0	0	1	0	0	0	8	
1931	0	0	0	0	0	0	7	1	0	0	0	0	0	0	0	0	0	0	0	8	
1932	0	0	0	0	9	0	13	1	0	0	0	0	0	0	0	0	0	0	0	23	
1933	0	0	0	0	8	0	13	1	0	0	0	0	0	0	0	0	0	0	0	22	
1934	0	0	0	1	21	0	20	1	0	0	0	0	0	0	0	0	0	0	0	43	
1935	0	0	0	9	9	0	20	1	0	0	0	0	0	0	0	1	0	0	0	40	
1936	0	0	0	12	14	0	15	0	0	0	0	0	0	0	0	0	0	0	0	41	
1937	0	0	0	13	17	0	37	0	0	0	0	0	0	0	0	1	0	0	0	68	
1938	0	0	0	15	20	0	44	0	0	0	0	0	0	0	0	0	1	0	0	80	
1939	0	0	0	18	24	0	44	1	0	0	0	0	0	2	0	0	0	0	0	89	
1940	0	0	0	15	33	0	52	0	0	0	0	0	0	0	0	0	1	0	0	101	
1941	0	0	0	40	40	0	37	1	0	0	0	0	0	2	0	0	0	0	0	120	
1942	0	0	0	53	67	0	44	0	0	0	0	0	0	1	0	0	1	0	0	166	
1943	0	0	0	42	51	0	67	1	0	0	0	0	0	0	0	0	0	0	0	161	
1944	0	0	0	38	47	0	52	0	0	0	0	0	0	0	0	0	1	0	0	138	
1945	0	0	0	3	2	0	44	0	0	0	0	0	0	0	0	0	0	0	0	49	
1946	0	0	0	11	21	14	51	4	0	0	0	0	0	1	0	0	4	0	0	106	
1947	0	0	0	19	21	27	57	7	0	0	0	0	0	0	0	0	8	0	0	139	
1948	0	3	0	22	26	56	57	1	0	0	1	0	0	0	0	0	26	0	0	192	
1949	0	0	0	25	31	20	61	0	0	0	1	0	2	0	0	5	6	0	2	0	153
1950	0	3	0	29	37	15	63	41	0	0	2	0	1	0	13	18	0	0	0	222	
1951	1	1	0	31	40	62	87	9	0	3	0	0	0	0	5	14	0	0	0	253	
1952	0	1	0	36	45	142	92	1	0	0	0	0	1	0	9	20	0	0	0	347	
1953	0	0	0	42	50	90	75	1	0	0	3	0	0	0	38	35	1	0	0	335	
1954	0	0	1	43	54	35	24	26	0	0	0	0	0	0	32	59	1	0	0	275	
1955	0	0	0	49	60	20	108	11	0	0	2	0	0	0	20	43	1	1	0	315	
1956	0	0	0	54	62	16	140	25	0	1	3	0	0	0	47	69	0	0	0	417	
1957	17	1	0	59	70	2	111	14	2	0	1	0	0	0	31	33	1	0	0	342	
1958	0	0	0	67	65	0	126	13	0	0	1	0	0	0	0	86	0	0	0	358	
1959	0	0	0	78	71	0	69	7	0	0	0	0	0	0	0	47	0	0	0	272	
1960	0	0	0	72	59	0	64	6	0	1	1	0	0	0	0	41	0	0	0	244	
1961	0	0	0	39	28	0	81	9	0	0	0	0	0	0	0	56	0	0	0	213	
1962	0	0	0	55	52	0	46	7	0	0	0	0	0	0	0	48	0	0	0	208	
1963	0	0	0	122	52	0	49	6	0	0	0	0	0	0	0	40	0	0	0	269	
1964	0	0	0	139	95	6	85	6	0	0	0	0	0	0	0	39	0	0	0	370	
1965	0	1	0	83	101	11	51	3	0	0	0	0	0	0	0	62	0	0	0	312	
1966	0	2	0	76	87	0	81	8	1	0	0	0	0	0	0	71	0	0	0	326	
1967	0	0	0	109	73	2	50	50	6	0	0	0	0	0	0	2	55	0	0	0	297
1968	0	0	0	98	75	8	58	4	1	0	0	0	0	0	2	0	22	0	0	0	268
1969	0	0	0	118	95	10	27	2	0	0	0	0	3	0	7	43	0	0	0	305	
1970	0	0	0	186	188	5	101	5	1	0	0	2	4	0	8	38	0	0	2	540	
1971	0	0	0	200	189	3	84	6	0	0	0	0	0	0	8	54	1	0	0	545	
1972	0	0	0	252	286	0	35	17	0	0	0	0	0	0	0	78	0	0	0	668	
1973	0	0	0	215	244	0	83	26	0	2	14	0	0	0	15	95	2	28	0	724	
1974	0	0	0	213	271	0	63	34	0	9	0	0	0	1	5	44	4	22	0	666	
1975	0	0	0	196	293	9	35	63	0	3	0	0	0	18	2	62	11	1	0	693	
1976	0	0	0	353	174	0	35	27	0	0	0	0	0	0	10	89	0	0	0	688	
1977	0	0	0	234	304	0	32	71	0	0	0	0	0	0	0	58	0	0	0	699	
1978	0	0	0	181	354	0	93	133	0	0	0	0	0	0	0	19	0	0	0	780	
1979	0	0	0	164	379	0	95	150	0	0	0	0	0	0	8	17	0	0	0	813	
1980	0	0	0	447	147	0	88	72	0	0	0	0	0	0	10	40	0	0	0	804	
1981	0	1	0	188	192	0	148	39	1	0	0	0	0	0	13	28	0	0	0	610	
1982	0	0	0	229	210	2	105	56	1	0	0	0	0	0	9	5	0	0	0	617	
1983	0	0	0	100	142	3	66	68	0	0	0	0	0	0	6	4	0	0	0	389	
1984	0	0	0	87	105	0	64	88	0	0	0	0	0	0	0	46	0	0	0	390	
1985	0	0	1	23	29	5	39	123	0	0	0	0	0	0	2	30	0	0	0	252	
1986	0	0	0	1	31	20	69	89	0	0	0	0	0	0	0	19	0	0	0	229	
1987	0	0	0	0	0	0	80	86	0	0	0	0	0	0	0	16	0	0	0	182	
1988	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1989	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1990	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1992	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1993	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
1994	0	0	0	0	0	0	0	0	0	0	0	0	18	0	0	0	0	0	0	0	
1995	0	0	0	0	0	0	0	0	0	0	0	91	0	0	0	0	0	0	0	91	
1996	0	0	0	0	0	0	0	28	0	0	16	0	0	0	0	19	0	0	0	63	
1997	0	0	0	0	0	0	0	0	1	1	30	55	0	0	0	0	0	0	0	87	
1998	0	0	0	0	0	0	0	0	22	26	41	0	0	0	0	0	0	0	0	89	
1999	0	0	0	0	0	0	2	39	2	0	0	0	0	0	0	28	0	0	0	71	

Table 2. Males contd.

	1E	2C	2R	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13	Total
2000	0	0	0	0	0	0	4	15	0	0	0	16	0	0	0	0	0	0	0	35
2001	0	0	0	0	0	0	11	10	19	7	20	26	0	0	0	0	0	0	0	93
2002	0	0	0	0	0	0	0	79	1	0	8	31	0	0	0	0	0	0	0	119
2003	0	0	0	0	0	0	32	0	4	7	35	37	0	0	0	0	0	0	0	115
2004	0	0	0	0	0	0	0	62	0	0	0	75	0	0	0	0	0	1	0	138
2005	0	0	0	0	0	0	28	67	2	0	7	52	0	0	0	0	0	0	0	156
2006	0	0	0	0	0	0	41	33	11	1	36	23	0	0	0	0	0	0	0	145
2007	0	0	0	0	0	0	50	67	3	0	15	5	0	0	0	0	0	0	0	140
2008	0	0	0	0	0	0	23	33	0	0	5	48	0	0	0	0	0	0	0	109
2009	0	0	0	0	0	0	29	41	8	3	13	6	0	0	0	0	0	0	0	100
2010	0	0	0	0	0	0	17	40	0	0	0	12	0	0	0	0	0	0	0	69
2011	0	0	0	0	0	0	17	64	0	0	0	1	0	0	0	0	0	0	0	82
2012	0	0	0	0	0	0	47	61	4	0	3	0	0	0	0	0	0	0	0	115
2013	0	0	0	0	0	0	17	41	0	0	0	3	0	0	0	0	0	0	0	61
2014	0	0	0	0	0	0	16	35	0	0	0	0	0	0	0	0	0	0	0	51
2015	0	0	0	0	0	0	10	35	0	0	0	0	0	0	0	0	0	0	0	45
2016	0	0	0	0	0	0	7	8	0	0	0	0	0	0	0	0	0	0	0	15
2017	0	0	0	0	0	0	3	22	6	10	4	17	0	0	0	0	9	0	0	71
2018	0	0	0	0	0	0	28	22	4	1	15	14	0	0	0	0	16	0	0	100
2019	0	0	0	0	0	0	26	32	3	0	0	0	0	0	0	0	5	0	0	66
2020	0	0	0	0	0	0	1	58	0	0	0	0	0	0	0	4	0	0	0	63
Total	18	13	2	5,004	5,270	583	3,953	2,270	97	75	277	532	17	21	305	1,775	22	55	2	20,291

Females

	1E	2C	2R	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13	Total	
1930	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	1	0	0	0	5	
1931	0	0	0	0	0	0	4	0	0	0	0	0	0	0	0	2	0	0	0	6	
1932	0	0	0	5	4	0	7	0	0	0	0	0	0	0	0	1	0	0	0	17	
1933	0	0	0	5	4	0	7	1	0	0	0	0	0	0	1	0	1	0	0	19	
1934	0	0	0	9	10	0	10	0	0	0	0	0	0	0	1	0	1	0	0	31	
1935	0	0	0	8	14	0	10	0	0	0	0	0	0	0	0	1	0	0	0	33	
1936	0	0	0	12	13	0	7	0	0	0	0	0	0	0	0	2	0	0	0	34	
1937	0	0	0	14	18	0	18	1	0	0	0	0	0	0	0	1	0	0	0	52	
1938	0	0	0	18	20	0	22	0	0	0	0	0	0	0	0	1	0	0	0	61	
1939	0	0	0	19	23	0	22	0	0	0	0	0	0	1	0	2	0	0	1	68	
1940	0	0	0	13	34	0	25	0	0	0	0	0	0	0	0	1	0	0	0	73	
1941	0	0	0	64	38	0	18	0	0	0	0	0	0	0	0	2	0	0	0	122	
1942	0	0	0	54	66	0	22	0	0	0	0	0	0	2	0	0	1	0	0	145	
1943	0	0	0	39	51	0	32	0	0	0	0	0	0	0	0	2	0	0	0	124	
1944	0	0	0	38	45	0	25	0	0	0	0	0	0	0	0	1	0	0	0	109	
1945	0	0	0	2	3	0	22	1	0	0	0	0	0	0	0	2	0	0	0	30	
1946	0	0	0	10	18	10	24	1	0	0	0	0	1	0	0	13	0	0	0	77	
1947	0	0	0	18	19	21	27	3	0	0	0	0	0	0	0	23	0	0	0	111	
1948	0	0	0	21	25	38	31	0	0	0	0	0	0	0	0	53	0	0	0	168	
1949	0	0	0	25	31	30	32	0	0	0	2	0	0	0	4	27	0	1	0	152	
1950	0	1	1	29	34	9	25	19	0	0	0	0	0	0	0	32	0	1	0	151	
1951	0	0	0	33	42	39	42	2	0	2	1	0	2	0	2	70	0	1	0	236	
1952	0	0	1	37	45	43	78	2	0	0	0	0	1	0	0	97	1	0	0	305	
1953	0	0	0	39	49	47	56	2	0	0	3	0	0	0	5	57	1	0	0	259	
1954	0	1	0	45	55	27	22	15	0	0	3	0	1	0	4	124	0	0	0	297	
1955	0	0	0	58	59	15	80	4	0	0	3	0	0	0	7	119	0	2	0	347	
1956	0	0	0	62	66	23	97	7	0	0	1	0	1	0	13	108	0	4	0	382	
1957	11	1	0	79	68	0	81	12	2	0	3	0	0	0	0	13	96	1	0	0	367
1958	0	0	0	101	63	0	128	8	0	0	1	0	0	0	0	153	0	0	0	454	
1959	0	0	0	126	73	0	70	4	0	0	0	0	0	0	0	83	0	1	0	357	
1960	0	0	0	141	57	0	65	4	0	1	1	0	0	0	0	73	0	0	0	342	
1961	0	0	0	82	30	0	83	5	0	0	1	0	0	0	0	98	0	0	0	299	
1962	0	0	0	117	52	0	47	5	0	0	0	0	0	0	0	85	0	1	0	307	
1963	0	0	0	168	52	0	50	4	0	0	0	0	0	0	0	71	0	0	0	345	
1964	0	0	0	186	97	6	86	4	0	0	0	0	0	0	0	69	0	0	0	448	
1965	0	1	0	110	102	9	99	3	0	0	0	0	0	0	0	94	0	0	0	418	
1966	0	1	0	105	88	2	100	15	0	0	0	0	0	0	0	84	0	0	0	395	
1967	0	0	0	139	73	8	65	7	0	0	0	0	0	0	0	3	87	0	0	0	382
1968	0	0	0	124	73	3	81	3	0	0	0	0	0	0	7	5	56	0	0	0	352
1969	0	0	0	156	96	10	32	1	0	0	0	0	0	8	0	5	97	0	0	0	405
1970	0	0	0	216	188	2	87	5	1	0	0	0	0	0	0	4	70	0	0	2	575
1971	0	0	0	250	190	2	67	4	0	0	0	0	0	0	0	9	52	0	0	0	574
1972	0	0	0	292	286	0	75	22	0	0	0	0	0	0	0	1	113	0	0	0	789
1973	0	0	0	239	244	2	90	15	0	2	7	0	0	0	0	6	116	11	27	0	759
1974	0	0	0	267	272	0	51	19	0	3	0	0	0	0	0	3	79	17	18	0	729
1975	0	0	0	229	288	2	46	22	0	4	0	0	0	0	2	4	58	23	0	0	678
1976	0	0	0	445	174	0	46	29	0	0	0	0	0	0	0	11	113	0	0	1	819

Table 2. Females contd.

	1E	2C	2R	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13	Total
1977	0	0	0	269	303	0	28	14	0	0	0	0	0	0	2	43	0	0	0	659
1978	0	0	0	207	356	0	85	22	0	0	0	0	0	0	0	48	0	0	0	718
1979	0	0	0	130	264	0	38	28	0	0	0	0	0	0	0	7	64	0	0	531
1980	0	0	0	272	109	0	70	12	0	0	0	0	0	0	0	5	82	0	0	550
1981	0	0	0	188	192	0	68	11	0	0	0	0	0	0	0	2	63	0	0	524
1982	0	0	0	236	219	2	58	28	0	0	0	0	0	0	0	6	56	0	0	605
1983	0	0	0	98	138	4	69	30	0	0	0	0	0	0	0	5	42	0	0	386
1984	0	0	0	87	114	0	38	55	0	0	0	0	0	0	0	0	76	0	0	370
1985	0	0	0	26	35	4	20	41	0	0	0	0	0	0	0	5	66	0	0	197
1986	0	0	0	0	15	2	35	43	2	0	0	0	0	0	0	0	54	0	0	151
1987	0	0	0	0	0	0	43	30	0	0	0	0	0	0	0	0	49	0	0	122
1988	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1989	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1990	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1991	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1992	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1993	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1994	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	3
1995	0	0	0	0	0	0	0	0	0	0	9	0	0	0	0	0	0	0	0	9
1996	0	0	0	0	0	0	0	2	1	0	0	0	0	0	0	0	11	0	0	14
1997	0	0	0	0	0	0	0	0	0	0	1	12	0	0	0	0	0	0	0	13
1998	0	0	0	0	0	0	0	0	3	4	4	0	0	0	0	0	0	0	0	11
1999	0	0	0	0	0	0	0	7	0	0	0	0	0	0	0	0	22	0	0	29
2000	0	0	0	0	0	0	1	4	0	0	0	0	0	0	0	0	0	0	0	5
2001	0	0	0	0	0	0	0	0	3	0	1	3	0	0	0	0	0	0	0	7
2002	0	0	0	0	0	0	0	31	0	0	0	2	0	0	0	0	0	0	0	33
2003	0	0	0	0	0	0	30	0	1	0	3	2	0	0	0	0	0	0	0	36
2004	0	0	0	0	0	0	0	14	0	0	0	8	0	0	0	0	0	0	0	22
2005	0	0	0	0	0	0	37	19	0	0	7	3	0	0	0	0	0	0	0	66
2006	0	0	0	0	0	0	35	12	1	1	2	1	0	0	0	0	0	0	0	52
2007	0	0	0	0	0	0	46	21	0	0	0	1	0	0	0	0	0	0	0	68
2008	0	0	0	0	0	0	38	18	0	0	0	6	0	0	0	0	0	0	0	62
2009	0	0	0	0	0	0	35	24	0	0	5	1	0	0	0	0	0	0	0	65
2010	0	0	0	0	0	0	28	20	0	0	0	2	0	0	0	0	0	0	0	50
2011	0	0	0	0	0	0	6	37	0	0	0	1	0	0	0	0	0	0	0	44
2012	0	0	0	0	0	0	38	30	1	0	0	0	0	0	0	0	0	0	0	69
2013	0	0	0	0	0	0	17	17	0	0	0	0	0	0	0	0	0	0	0	34
2014	0	0	0	0	0	0	14	16	0	0	0	0	0	0	0	0	0	0	0	30
2015	0	0	0	0	0	0	9	16	0	0	0	0	0	0	0	0	0	0	0	25
2016	0	0	0	0	0	0	9	13	0	0	0	0	0	0	0	0	0	0	0	22
2017	0	0	0	0	0	0	0	13	0	1	0	6	0	0	0	0	38	0	0	58
2018	0	0	0	0	0	0	23	8	0	0	1	8	0	0	0	0	31	0	0	71
2019	0	0	0	0	0	0	20	10	0	0	0	0	0	0	0	0	27	0	0	57
2020	0	0	0	0	0	0	5	25	0	0	0	0	0	0	0	2	0	0	0	32
Total	11	5	2	5,762	5,097	360	3,161	920	15	18	50	68	17	11	131	3,266	54	56	4	19,008

Table 3

The High Catch Series.

Catches for the years and sub-areas where they differ from the 'best' catch series (1930-1, 1936-45 in sub-areas 7CS, 7CN and 11; 1947-56 in sub-areas 5 and 6W). Numbers from the 'best' catch series are shown for comparison. The 'high' catch series is identical to the 'best' series for all other areas and years.

Series: Sub-area:	Best 7CS	Best 7CS	High 7CS	High 7CS	Best 7CN	Best 7CN	High 7CN	High 7CN	Best 11	Best 11	High 11	High 11
	Male	Fem	Male	Fem			Male	Fem			Male	Fem
1930	7	4	14	8	0	0	0	0	1	1	2	2
1931	7	4	14	8	1	0	2	0	0	2	0	4
1932	13	7	13	7	1	0	1	0	0	1	0	1
1933	13	7	13	7	1	1	1	1	0	1	0	1
1934	20	10	20	10	1	0	1	0	0	1	0	1
1935	20	10	20	10	1	0	1	0	1	1	1	1
1936	15	7	30	14	0	0	0	0	0	2	0	4
1937	37	18	74	36	0	1	0	2	1	1	2	2
1938	44	22	88	44	0	0	0	0	1	1	2	2
1939	44	22	88	44	1	0	2	0	0	2	0	4
1940	52	25	104	50	0	0	0	0	1	1	2	2
1941	37	18	74	36	1	0	2	0	0	2	0	4
1942	44	22	88	44	0	0	0	0	1	1	2	2
1943	67	32	134	64	1	0	2	0	0	2	0	4
1944	52	25	104	50	0	0	0	0	1	1	2	2
1945	44	22	44	22	0	1	0	2	0	2	0	4

Table 3 continued

Series: Sub-area:	Best 5	Best 5	High 5	High 5	Best 6W	Best 6W	High 6W	High 6W
	Male	Fem	Male	Fem	Male	Fem	Male	Fem
1946	11	10	11	10	21	18	21	18
1947	19	18	55	56	21	19	70	68
1948	22	21	55	56	26	25	70	68
1949	25	25	55	56	31	31	70	68
1950	29	29	55	56	37	34	70	68
1951	31	33	55	56	40	42	70	68
1952	36	37	55	56	45	45	70	68
1953	42	39	55	56	50	49	70	68
1954	43	45	55	56	54	55	70	68
1955	49	58	56	66	60	59	70	68
1956	54	62	57	66	62	66	70	68
1957	59	79	59	79	70	68	70	68

Table 4

Catch series for Trials 8 and 9 used to test the sensitivity to the allocation of catches off Korea between sub-areas 5 and 6W. Catches in the other sub-areas are the same as for the 'Best' catch series.

Sub-area:	Trial 8				Trial 9			
	5	5	6W	6W	5	5	6W	6W
	Male	Fem	Male	Fem	Male	Fem	Male	Fem
1932	0	5	9	4	0	5	9	4
1933	0	5	8	4	0	5	8	4
1934	1	9	21	10	1	9	21	10
1935	9	12	9	10	7	7	12	14
1936	14	15	13	9	9	10	15	17
1937	17	16	14	15	12	9	21	20
1938	19	22	16	16	14	13	24	22
1939	23	23	20	18	15	15	27	27
1940	21	21	27	26	12	11	37	35
1941	48	72	31	31	38	62	41	41
1942	66	66	53	55	43	43	77	77
1943	51	51	40	41	31	33	59	60
1944	48	48	37	35	31	31	53	53
1945	3	2	2	3	3	2	2	3
1946	14	15	15	16	10	8	22	20
1947	24	21	16	16	15	15	23	24
1948	27	26	20	21	18	18	28	30
1949	30	32	25	25	18	22	36	36
1950	34	38	28	29	23	24	42	40
1951	40	40	33	33	26	26	47	47
1952	46	46	37	34	29	30	51	53
1953	50	51	40	39	31	33	58	58
1954	55	54	43	45	35	35	64	63
1955	62	69	46	49	39	48	70	69
1956	67	74	52	51	42	53	75	74
1957	73	92	56	55	49	66	79	82
1958	80	114	51	51	53	89	77	77
1959	93	141	57	57	63	110	86	89
1960	84	152	46	47	63	131	68	67
1961	44	87	24	24	35	77	33	34
1962	65	128	43	40	49	110	58	59
1963	131	179	43	41	104	149	71	70
1964	159	205	77	76	118	162	119	118
1965	102	131	82	81	68	97	116	115
1966	95	121	70	70	64	91	100	101
1967	125	153	59	57	91	120	93	90
1968	112	139	60	59	82	107	91	90
1969	137	176	75	77	98	138	114	115
1970	223	253	151	151	152	183	221	222
1971	239	286	152	152	165	214	225	225
1972	308	348	229	231	230	267	311	308
1973	251	275	208	208	197	220	262	263
1974	251	302	235	235	188	241	297	297
1975	253	287	235	231	159	196	327	324
1976	389	479	139	139	292	384	235	235
1977	294	331	242	243	192	226	346	346
1978	253	276	283	286	152	175	384	387
1979	164	130	379	264	164	130	379	264
1980	447	272	147	109	447	272	147	109
1981	188	188	192	192	188	188	192	192
1982	236	247	202	209	222	229	217	226
1983	100	98	142	138	100	98	142	138
1984	87	87	105	114	87	87	105	114
1985	23	26	29	35	23	26	29	35
1986	1	0	31	15	1	0	31	15

Bycatches

Tables 5 and 6 summarise recent bycatches (also referred to as incidental catches) off Japan and Korea by sub-area. Individual records, including position, date, length and sex, have been provided to the IWC by Japan for 1,964 by caught minke whales from 2001-16 received 28 May 2019) and by Korea for 1,883 by caught and stranded minke whales from 2001-17 (received 29 Mar 2019).

Table 5

Recent bycatches by Japan and Korea (some are updates to those listed in progress reports). It is known that the numbers off Japan in 2001 are incomplete. Bycatches from sub-area 6W by Japan are included with those in 6E (see text). No data for 2020 off Korea are available so the average catch over the previous 10 years is used. Bycatches that are shown in grey are not used in the fitting process.

Year	Japan							Korea					
	1E	2C	6E	7CN	7CS	10E	11	Total	5	6W	1W	Posn.Unk	Total
1996									0	128	0	0	128
1997									0	78	0	0	78
1998									0	47	0	0	47
1999									0	54	0	0	54
2000									12	80	0	0	92
2001	1	10	25	3	8	4	3	54	9	141	0	0	150
2002	7	19	45	13	17	3	5	109	8	75	0	0	83
2003	5	17	61	15	18		8	124	10	75	2	0	87
2004	4	19	66	9	14		3	115	9	52	0	0	61
2005	4	33	55	10	17	3	6	128	7	98	0	0	105
2006	3	28	76	16	21		3	147	11	67	0	2	80
2007	7	42	69	11	20		6	155	12	59	0	1	72
2008	9	23	68	11	17	2	3	133	12	61	0	2	75
2009	3	17	69	3	25		1	118	10	70	0	2	82
2010	3	18	74	8	17		4	124	8	63	0	0	71
2011	6	28	65	9	8		1	117	15	70	0	1	86
2012	5	25	56	9	15		4	114	8	66	0	0	74
2013	5	20	54	9	15	2		105	8	43	0	0	51
2014	3	21	74	16	23	1	2	140	7	43	0	0	50
2015	5	28	84	12	26		1	156	7	78	1	1	87
2016	7	34	86	17	22	3		169	10	84	0	0	94
2017	5	32	80	10	34	1	2	164	12	57	0	0	69
2018	2	18	40	9	18			87	7	73	0	0	80
2019	3	15	54	9	23			104	3	55	0	0	58
2020	2	10	34	9	16			71	[9]	[63]	0	0	[96]
Total	89	457	1235	374	208	19	52	2434					

In Japan it has been obligatory to report bycatches from 2001, since when the numbers are considered to be reliable. Almost all of the reported bycatch off Japan occurs in set-net fisheries. Three types of set nets are used: large-scale (excluding salmon nets), salmon nets and small-scale. For fishing gears other than set-nets, bycatch, retention and marketing of whales are prohibited by the 2001 regulation and a diagnostic DNA registry is used to deter illegal distribution of any whales caught.

Ideally, the catch by each gear type should be modelled separately to allow the historical (pre-2001) bycatch to be predicted. However, information on numbers of catches by net type is not available. Therefore, in the 2013 Implementation, the historical bycatches for each sub-area were set using the total number of bycatches and the combined number of large-scale and salmon nets in each sub-area (see Allison, 2014). Japan has provided new information on the numbers of large-scale nets from 1979-2018, but numbers of salmon nets since 2016 are not available. The numbers of whales caught in salmon nets are small in comparison to those from large-scale nets (see Table 6 which lists the bycatches from 2001-9 by gear type), so in the current implementation, the historical bycatches are set using the total numbers of bycatches (Table 5) and the numbers of large-scale nets (Table 7) in each sub-area.

Table 6

Bycatches by Japan 2001-9 by gear type (Hakamada pers comm 28/04/2011).

	Salmon	Large-scale	Small-scale	Other gear types
2001	3	68	9	0
2002	5	92	12	0
2003	8	99	18	0
2004	2	101	12	2
2005	7	105	17	0
2006	5	125	17	0
2007	8	131	16	0
2008	3	116	14	0
2009	4	101	13	0

Table 7
Numbers of nets.

	Japan large-scale trap nets								Korean nets		
	1E	2C	6E	7CS	7CN	10E	11	Total	5	6W	Total
1946	24	67	103	41	8	7	3	252	0	0	0
1947	26	73	112	44	9	7	3	275	2	5	7
1948	29	79	122	48	9	8	4	298	4	11	15
1949	31	85	131	52	10	8	4	321	6	16	22
1950	33	91	141	55	11	9	4	344	8	21	29
1951	35	97	150	59	11	10	5	367	10	27	37
1952	37	103	159	63	12	10	5	390	12	32	44
1953	40	109	169	66	13	11	5	412	14	38	52
1954	42	115	178	70	14	11	5	435	15	43	58
1955	44	121	187	74	14	12	6	458	17	48	65
1956	46	127	197	77	15	13	6	481	19	54	73
1957	48	133	206	81	16	13	6	504	21	59	80
1958	51	139	216	85	16	14	7	527	23	64	87
1959	53	145	225	88	17	14	7	550	25	70	95
1960	55	151	234	92	18	15	7	573	27	75	102
1961	57	157	244	96	19	16	7	596	29	80	109
1962	59	164	253	100	19	16	8	619	31	86	117
1963	62	170	262	103	20	17	8	642	33	91	124
1964	64	176	272	107	21	17	8	665	35	97	132
1965	66	182	281	111	21	18	9	687	37	102	139
1966	68	188	291	114	22	19	9	710	39	107	146
1967	70	194	300	118	23	19	9	733	41	113	154
1968	73	200	309	122	24	20	9	756	43	118	161
1969	75	206	319	125	24	20	10	779	44	123	167
1970	77	212	328	129	25	21	10	802	46	129	175
1971	80	209	324	127	25	21	10	795	48	134	182
1972	83	206	321	124	25	21	10	789	50	139	189
1973	86	203	317	122	24	20	9	782	52	145	197
1974	89	200	314	119	24	20	9	776	54	150	204
1975	92	197	310	117	24	20	9	769	56	156	212
1976	80	198	321	118	25	21	10	773	58	161	219
1977	69	199	332	119	27	22	10	777	60	166	226
1978	57	200	344	119	28	23	11	781	62	172	234
1979	46	205	361	122	30	25	11	800	64	177	241
1980	49	208	372	130	28	24	11	822	66	182	248
1981	51	205	375	134	26	21	10	823	68	188	256
1982	50	204	393	133	27	22	10	838	70	193	263
1983	54	199	392	132	37	31	14	859	71	198	269
1984	51	191	393	141	48	41	19	885	73	204	277
1985	47	192	419	141	42	36	16	894	75	209	284
1986	50	198	413	136	50	43	20	909	77	215	292
1987	47	196	409	138	48	41	19	900	79	220	299
1988	47	190	407	132	40	33	15	865	81	225	306
1989	56	185	398	142	35	29	13	857	83	231	314
1990	56	182	413	137	35	30	14	867	85	236	321
1991	61	178	410	135	29	24	11	847	85	286	371
1992	56	169	400	135	27	22	10	820	96	305	401
1993	62	183	406	135	28	22	10	845	96	291	387
1994	55	179	387	131	29	23	11	814	94	286	380
1995	57	179	380	119	26	20	9	790	97	292	389
1996	57	175	379	132	26	20	9	799	103	352	455
1997	54	172	376	132	25	19	9	787	123	340	463
1998	56	167	377	133	26	19	9	787	105	338	443
1999	55	170	370	131	28	22	10	786	120	321	441
2000	55	169	367	130	28	22	10	781	105	318	423
2001	58	154	367	132	29	23	11	775	82	311	393
2002	52	163	367	130	32	26	12	781	88	292	380
2003	49	164	363	137	31	25	12	781	81	286	367
2004	51	160	351	137	27	21	10	757	94	267	361
2005	52	159	329	132	26	21	9	729	81	263	344
2006	45	156	313	132	27	21	10	703	78	255	333
2007	43	146	329	123	8	4	2	654	77	247	324
2008	40	129	315	121	22	16	7	651	71	230	301
2009	42	130	311	121	22	16	7	648	68	219	287
2010	40	130	314	116	21	15	7	644	68	219	287
2011	40	130	311	94	21	15	7	617	68	219	287
2012	39	128	313	96	20	14	7	617	68	219	287
2013	38	120	307	92	20	14	7	598	68	219	287
2014	36	120	300	97	20	15	7	594	68	219	287
2015	35	115	300	101	20	15	7	592	68	219	287
2016	37	120	280	101	21	15	7	582	68	219	287
2017	36	122	275	93	21	15	7	569	68	219	287
2018	34	114	288	87	21	15	7	567	68	219	287
2019	34	114	288	87	21	15	7	567	68	219	287
2020	34	114	288	87	21	15	7	567	68	219	287

Table 7 contd.

Sources: Japan 1935-70. Set using linear interpolation, assuming 0 in 1935.
Japan 1970-79. Set using linear interpolation between the numbers for 1970 and 1975 from Tobayama *et al.* (1992).
Japan 1979-2018. Pastene, pers. comm. Apr 2021.
Korea 1946-1996. Set using linear interpolation, assuming 0 in 1946.
Korea 1990-2009. An, pers. comm.

Missing data: where the numbers of nets are unknown (off Japan from 2019-20 and off Korea from 2010-20), the last known numbers are used.

The bycatch in sub-area 6W by Japan is small (9 whales) (and there are no corresponding set net numbers) so the numbers are added to the bycatches for sub-area 6E. The bycatch by Korea in sub-area 1W is very small (3 whales in total) and there are no corresponding set net numbers so the numbers are added to the bycatches for sub-area 5.

Japan updated the numbers of large-scale nets up to and including 2018 and incorporated information from the Japanese Coast Guard for 2014 on the dates that the nets were in operation (see IWC 2020). The set nets are assigned to sub-area based on the position of the centre of the net, although some nets extend beyond a single sub-area. Korea provided revised data on the number of set nets in operation based on the number of licenses issued between 1994-2017 (see IWC 2020), but the Committee (Item 8.1.3) decided that the number of nets provide a more reliable source of information than the number of licenses.

For the best effort series, the numbers of nets off Japan are extrapolated from 1946 to 1969 assuming a linear relationship from 0 in 1935 to the known numbers in 1970 (Tobayama *et al.*, 1992). Bycatches before 1946 are ignored because, although some set-nets were in operation before 1946 (Brownell, pers. comm.), the numbers are highly uncertain and are sufficiently small that they are unlikely to affect the implementation.

The numbers of nets are listed in Table 7. The numbers of bycatches are only used in the trials where the number of nets is also known. Thus, for example for Japan, the catches from 2019-20 are not used and are shown greyed out in table 5.

A single series of historical bycatches is used for all of the trials when testing the effect of future catches (including those set by the RMP), irrespective of the true values of the bycatches, which differ both among trials and simulations within trials. The estimate of the future bycatch is set to the averages of the predicted bycatches based on the fit to the actual data of the operating model for the six baseline trials (i.e. using the 'best fit' simulation (0)). This series will be generated once conditioning is complete.

A sensitivity trial will be undertaken that assumes catches since 2001 were under-reported by 50% and the numbers of nets were double the best-case values from 1946-1969 (up to a maximum equal to the number of nets in 1969).

Korea: The same method as used for Japan is applied, except that bycatch numbers since 1945 are extrapolated from reported numbers in sub-areas 5 (Yellow Sea) and 6W (East Sea) since 2000 and 1996 respectively. Catches in sub-area 6W are assumed to be under-reported by 50% (based on DNA profiling and a capture-recapture analysis of market products, Baker *et al.*, 2007). The high effort sensitivity trial assumes that bycatches in sub-area 5 since 2000 were under-reported by 50%, and the numbers of nets were double the best-case values from 1946-1969 (up to a maximum equal to the number of nets in 1969).

China: There are no data on bycatches off China, although they are known to occur. There are not many set-nets in operation off China and the operations are likely to be similar to those off western Korea. In the absence of information the baseline trials assume that the bycatch off China is double that off western Korea. A sensitivity trial ignores any possible bycatch off China.

References

- Allison, C. 2011. Direct catch data for western North Pacific minke whale simulation trials. Paper SC/D11/NPM3 presented to the First Intersessional Workshop for the Implementation Review of western North Pacific common minke whales, 12-16 December 2011, Tokyo, Japan (unpublished). [Paper available from the Office of this Journal].
- Allison, C. 2014. The Historical Catch Series. *J. Cetacean Res. Manage. (Suppl.)* 15: 151-7.
- Allison, C. 2020. IWC individual catch database Version 7.1 released 23 December 2020. Available from the International Whaling Commission, 135 Station Road, Impington, Cambridge, CB24 9NP UK. [Statistics@iwc.int]
- Gong, Y. 1982. A note on the distribution of minke whales in Korean waters. *Rep. int. Whal. Commn* 32:279-82. SC/33/Mi7.
- Hakamada, T. 2010. The number of set nets in the coast of Japan by sub-areas and years during 1979-2006. Paper SC/D10/NPM13 presented to the First Intersessional Workshop for Western North Pacific Common Minke Whales, 14-17 December 2010, Pusan, Republic of Korea (unpublished). 1pp. [Paper available from the Office of this Journal].
- International Whaling Commission. 2020. Report of the 'First' Intersessional Workshop on the Implementation Review for western North Pacific Common Minke Whales, 25 February-1 March 2019, Tokyo, Japan, Annex F Effort Information. *J. Cetacean Res. Manage. (Suppl.)* 21:393-4.
- Park, J.-Y. 1995. Whaling History in Korean Waters. 593pp. [In Korean].
- Tobayama, T., Yanagisawa, F., and Kasuya, T. 1992. Incidental Take of Minke Whales in Japanese Trap Nets. *Rep. Int. Whal. Commn* 42:433-436.
- Ohsumi, S. 1982 Minke whales in the coastal waters of Japan, 1980 and a population assessment of the Okhotsk Sea-West Pacific stock. *Rep. Int. Whal. Commn* 32, 1982, pp283-6. SC/33/Mi8.

Appendix 2

Using the Genetic Stock Assignment by Sub-Area to Inform the Mixing Matrices of the North Pacific Minke Whale *Implementation Simulation Trials*

C.L. de Moor, C. Allison, A.E. Punt

This appendix details the stock assignment by sub-area and sex used to develop the data used to estimate mixing matrices for the North Pacific minke whale *Implementation Simulation Trials*. The baseline mixing matrices for Hypothesis E were newly developed for these *Implementation Simulation Trials*, largely informed by the genetic assignment tables below. The baseline mixing matrices for Hypotheses A and B were only changed from those used during the 2013 *Implementation Simulation Trials* where the genetic assignment tables below strongly supported such changes.

Baseline Trials, Hypotheses A and B

For the baseline trials, the stock assignment for Hypotheses A and B is based on the “stock90” assignment by STRUCTURE in *Data_NPM_190226_v3.csv*. The number of samples assigned to stock by sub-area is as follows. Table 7a of specifications details the assigned numbers by stock, sub-area, period and sex used to condition the trials.

Males	10E	11	1E	2C	6E	7CN	7CS	7E	7WR	8	9
J-stock	8	28	29	107	453	158	135	0	0	0	1
O-stock	1	29	1	26	1	580	281	41	74	207	442
Unassigned	2	7	2	10	41	80	61	3	6	22	44
Females											
J-stock	6	28	42	188	471	112	151	0	1	0	0
O-stock	0	30	0	24	3	263	286	4	8	17	49
Unassigned	1	7	2	17	33	23	49	1	0	6	5

Grey highlight: stock **has** been assigned to a sub-area, but is not modelled in that sub-area in the mixing matrices

- The singleton assignment of a J-stock female to sub-area 7WR is ignored for the baseline trials, but in Trial 5 J-stock animals are assumed to be found in both sub-areas 7E and 7WR.
 - The singleton assignment of an O-stock male to sub-area 1E is ignored for modelling purposes
 - The singleton assignment of a J-stock male to sub-area 9 is small compared to the total sample size, and is therefore ignored for the baseline, but in Trial 5 J-stock animals are assumed to be found in sub-areas 8 and 9
 - The assignment of O-stock animals to sub-area 6E are very small compared to the total sample size, and O-stock animals are therefore not modelled to be found in sub-area 6E.

Pink highlight: females of a stock have not been assigned to a sub-area, but are modelled in that sub-area in the mixing matrices

- The sample sizes in sub-area 10E are low and one cannot therefore discount the presence of O-stock females in sub-area 10E.

Hypothesis A Baseline

J-Stock Baseline A (Matrix J-A)

Hypothesis A Baseline (contd.)

O-Stock Baseline A (Matrix O-AB) Blue indicates changes since 2013 ISTs.

Age/ Sex	Mon	Sub - Area																				
		1W	1E	2C	2R	3	4	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE
Juv	J-M			γ_{13}	4	4	4				4	γ_{16}							γ_{30}			
	Apr			γ_{14}	2	2	2				8	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}			$2\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	May			γ_{14}	2	2	2				8	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}		$2\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	Jun			γ_{14}	2	2	2				4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}		$4\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	Jul			γ_{15}	2	2	2				4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}		$4\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	Aug			γ_{15}	2	2	2				4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}		$4\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	Sep			γ_{15}	2	2	2				4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}		$4\gamma_{30}$	γ_{22}	γ_{23}	γ_{24}
	O-D			γ_{15}	4	4	4				4	$2\gamma_{16}$							$2\gamma_{30}$			
Ad.M	J-M			γ_{13}	4	4	4				1	γ_{16}							γ_{30}			
	Apr			γ_{14}	2	2	2				2	$2\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$			$2\gamma_{30}$	γ_{22}	γ_{23}	$3\gamma_{24}$
	May										2	$2\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$		$2\gamma_{30}$	γ_{22}	γ_{23}	$6\gamma_{24}$
	Jun										2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$		$4\gamma_{30}$	γ_{22}	γ_{23}	$6\gamma_{24}$
	Jul										2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$		$4\gamma_{30}$	γ_{22}	γ_{23}	$6\gamma_{24}$
	Aug										2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$		$4\gamma_{30}$	γ_{22}	γ_{23}	$6\gamma_{24}$
	Sep										2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	γ_{21}		$4\gamma_{30}$	γ_{22}	γ_{23}	$3\gamma_{24}$
	O-D			γ_{15}	4	4	4				1	γ_{16}							γ_{30}			
Ad.F	J-M			γ_{13}	4	4	4				1	γ_{16}							γ_{30}			
	Apr			γ_{14}	2	2	2				1	γ_{16}	$2\gamma_{17}$	$2\gamma_{18}$	$2\gamma_{19}$	$2\gamma_{20}$			γ_{30}	γ_{22}	γ_{23}	$3\gamma_{24}$
	May										1	γ_{16}	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$		γ_{30}	$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$
	Jun										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$		$2\gamma_{30}$	$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$
	Jul										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$		$2\gamma_{30}$	$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$
	Aug										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$		$2\gamma_{30}$	$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$
	Sep										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$2\gamma_{21}$		$2\gamma_{30}$	$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$
	O-D			γ_{15}	4	4	4				1	γ_{16}							γ_{30}			

Hypothesis B Baseline

Y-Stock Baseline B (Matrix Y-BE)

Hypothesis B Baseline (contd.)

J-Stock Baseline B (Matrix J-BE)

Age/ Sex	Mon	Sub - Area																				
		1W	1E	2C	2R	3	4	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE
Juv	J-M	2	2						2	4γ ₂₉	2γ ₁	2γ ₄						γ ₆	γ ₇			
	Apr	2	2						2	4γ ₂₉	2γ ₁	2γ ₄						γ ₆	γ ₇	2γ ₈	2γ ₈	
	May	2	2						2	4γ ₂₉	2γ ₂	2γ ₄						γ ₆	γ ₇	2γ ₈	2γ ₈	
	Jun	2	2						2	4γ ₂₉	2γ ₃	2γ ₄						γ ₆	γ ₇	2γ ₉	2γ ₉	
	Jul	2	2						2	4γ ₂₉	2γ ₃	2γ ₅						γ ₆	γ ₇	2γ ₉	2γ ₉	
	Aug	2	2						2	4γ ₂₉	2γ ₃	2γ ₅						γ ₆	γ ₇	2γ ₉	2γ ₉	
	Sep	2	2						2	4γ ₂₉	2γ ₃	2γ ₅						γ ₆	γ ₇	2γ ₉	2γ ₉	
	O-D	2	2						2	4γ ₂₉	2γ ₃	2γ ₅						γ ₆	γ ₇	2γ ₉		
Ad.M	J-M	2	1						4	4γ ₂₉	2γ ₁	2γ ₄						γ ₆	γ ₇			
	Apr	1							2	2γ ₂₉	4γ ₁	2γ ₄						γ ₆	2γ ₇	γ ₈	γ ₈	
	May	1							2	2γ ₂₉	4γ ₂	2γ ₄						2γ ₆	2γ ₇	γ ₈	2γ ₈	
	Jun	1							2	2γ ₂₉	2γ ₃	4γ ₄						2γ ₆	2γ ₇	γ ₉	2γ ₉	
	Jul	1							2	2γ ₂₉	2γ ₃	4γ ₅						γ ₆	γ ₇	γ ₉	2γ ₉	
	Aug	1							2	2γ ₂₉	2γ ₃	4γ ₅						γ ₆	γ ₇	γ ₉	2γ ₉	
	Sep	2	1						4	4γ ₂₉	2γ ₃	4γ ₅						γ ₆	γ ₇			
	O-D	4	1						2		2γ ₃	2γ ₅										
Ad.F	J-M	2	1						4	4γ ₂₉	γ ₁	γ ₄						γ ₆	γ ₇			
	Apr	1							2	2γ ₂₉	2γ ₁	γ ₄						2γ ₆	2γ ₇	γ ₁₀	γ ₁₀	
	May	1							2	2γ ₂₉	2γ ₂	γ ₄						2γ ₆	2γ ₇	γ ₁₁	2γ ₁₁	
	Jun	1							2	2γ ₂₉	γ ₃	γ ₄						2γ ₆	2γ ₇	γ ₁₂	2γ ₁₂	
	Jul	1							2	2γ ₂₉	γ ₃	γ ₅						γ ₆	γ ₇	γ ₁₂	2γ ₁₂	
	Aug	1							2	2γ ₂₉	γ ₃	γ ₅						γ ₆	γ ₇	γ ₁₂	2γ ₁₂	
	Sep	2	1						4	4γ ₂₉	γ ₃	γ ₅						γ ₆	γ ₇			
	O-D	4	1						2		γ ₃	γ ₅										

O-Stock Baseline A (Matrix O-AB) Blue indicates changes since 2013 ISTs.

Age/ Sex	Mon	Sub - Area																					
		1W	1E	2C	2R	3	4	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13
Juv	J-M	γ ₁₃	4	4	4					4		γ ₁₆							γ ₃₀				
	Apr	γ ₁₄	2	2	2					8	2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀				2γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	May	γ ₁₄	2	2	2					8	2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	γ ₂₁			2γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	Jun	γ ₁₄	2	2	2					4	4γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	Jul	γ ₁₅	2	2	2					4	4γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	Aug	γ ₁₅	2	2	2					4	4γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	Sep	γ ₁₅	2	2	2					4	4γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	γ ₂₄	
	O-D	γ ₁₅	4	4	4					4	2γ ₁₆							2γ ₃₀					
Ad.M	J-M	γ ₁₃	4	4	4					1		γ ₁₆							γ ₃₀				
	Apr	γ ₁₄	2	2	2					2	2γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀				2γ ₃₀	γ ₂₂	γ ₂₃	3γ ₂₄	
	May									2	2γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀	2γ ₂₁			2γ ₃₀	γ ₂₂	γ ₂₃	6γ ₂₄	
	Jun									2	4γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀	2γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	6γ ₂₄	
	Jul									2	4γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀	2γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	6γ ₂₄	
	Aug									2	4γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀	2γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	6γ ₂₄	
	Sep									2	4γ ₁₆	4γ ₁₇	4γ ₁₈	4γ ₁₉	4γ ₂₀	2γ ₂₁			4γ ₃₀	γ ₂₂	γ ₂₃	6γ ₂₄	
	O-D	γ ₁₅	4	4	4					1		γ ₁₆							γ ₃₀				
Ad.F	J-M	γ ₁₃	4	4	4					1		γ ₁₆							γ ₃₀				
	Apr	γ ₁₄	2	2	2					1		γ ₁₆	2γ ₁₇	2γ ₁₈	2γ ₁₉	2γ ₂₀			γ ₃₀	γ ₂₂	γ ₂₃	3γ ₂₄	
	May									1		γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	4γ ₂₁		γ ₃₀	2γ ₂₂	2γ ₂₃	9γ ₂₄	
	Jun									1		2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	4γ ₂₁		2γ ₃₀	2γ ₂₂	2γ ₂₃	9γ ₂₄	
	Jul									1		2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	4γ ₂₁		2γ ₃₀	2γ ₂₂	2γ ₂₃	9γ ₂₄	
	Aug									1		2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	4γ ₂₁		2γ ₃₀	2γ ₂₂	2γ ₂₃	9γ ₂₄	
	Sep									1		2γ ₁₆	γ ₁₇	γ ₁₈	γ ₁₉	γ ₂₀	2γ ₂₁		2γ ₃₀	2γ ₂₂	2γ ₂₃	3γ ₂₄	
	O-D	γ ₁₅	4	4	4					1		γ ₁₆							γ ₃₀				

Baseline Trials, Hypothesis E

For the baseline trials, stock assignment for Hypothesis E is based on the “geneland.stock2” assignment by GENELAND in *Data_NPM_190226_v3.csv*. The number of samples assigned to stock by sub-area is as follows. Table 7a of specifications details the assigned numbers by stock, sub-area, period and sex used to condition the trials.

Males	10E	11	1E	2C	6E	7CN	7CS	7E	7WR	8	9
J-stock	8	13	31	88	492	20	0	0	0	0	0
P-stock	0	39	0	10	0	384	217	0	0	0	0
O-stock	0	1	0	0	0	280	83	41	70	207	464
Unassigned	0	6	0	19	0	55	105	0	0	0	0
Females											
J-stock	7	18	44	156	500	17	0	0	0	0	0
P-stock	0	24	0	10	0	216	296	0	0	0	0
O-stock	0	4	0	0	0	54	18	5	7	22	49
Unassigned	0	17	0	26	0	75	118	0	0	0	0

Pink highlight: animals of a stock **have not** been assigned to a sub-area, but are modelled in that sub-area in the mixing matrices

- It is assumed the J-stock occurs distributed in sub-area 7CS given they have been assigned to sub-areas 7CN and 2C to the east of Japan as well as sub-areas 6E and 10E to the west of Japan.

Hypothesis E Baseline

Y-Stock Baseline E (Matrix Y-BE)

Hypothesis E Baseline (contd.)

J-Stock Baseline E (Matrix J-BE)

P-Stock Baseline E (Matrix P-E)

Hypothesis E Baseline (contd.)

O-Stock Baseline E (Matrix O-E)

Age/ Sex	Mon	Sub - Area																						
		1W	1E	2C	2R	3	4	5	6W	6E	7CS	7CN	7WR	7E	8	9	9N	10W	10E	11	12SW	12NE	13	
Juv	J-M					4	4	4			4	γ_{16}												
	Apr					2	2	2			8	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}					γ_{22}	γ_{23}	γ_{24}	
	May					2	2	2			8	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}				γ_{22}	γ_{23}	γ_{24}	
	Jun					2	2	2			4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}				γ_{22}	γ_{23}	γ_{24}	
	Jul					2	2	2			4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}				γ_{22}	γ_{23}	γ_{24}	
	Aug					2	2	2			4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}				γ_{22}	γ_{23}	γ_{24}	
	Sep					2	2	2			4	$4\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	γ_{21}				γ_{22}	γ_{23}	γ_{24}	
	O-D					4	4	4			4	$2\gamma_{16}$												
Ad.M	J-M					4	4	4			1	γ_{16}												
	Apr					2	2	2			2	$2\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$					γ_{22}	γ_{23}	$3\gamma_{24}$	
	May					0	0	0			2	$2\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$				γ_{22}	γ_{23}	$6\gamma_{24}$	
	Jun					0	0	0			2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$				γ_{22}	γ_{23}	$6\gamma_{24}$	
	Jul					0	0	0			2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$				γ_{22}	γ_{23}	$6\gamma_{24}$	
	Aug					0	0	0			2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	$2\gamma_{21}$				γ_{22}	γ_{23}	$6\gamma_{24}$	
	Sep					0	0	0			2	$4\gamma_{16}$	$4\gamma_{17}$	$4\gamma_{18}$	$4\gamma_{19}$	$4\gamma_{20}$	γ_{21}				γ_{22}	γ_{23}	$3\gamma_{24}$	
	O-D					4	4	4			1	γ_{16}												
Ad.F	J-M					4	4	4			1	γ_{16}												
	Apr					2	2	2			1	γ_{16}	$2\gamma_{17}$	$2\gamma_{18}$	$2\gamma_{19}$	$2\gamma_{20}$					γ_{22}	γ_{23}	$3\gamma_{24}$	
	May										1	γ_{16}	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$				$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$	
	Jun										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$				$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$	
	Jul										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$				$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$	
	Aug										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$4\gamma_{21}$				$2\gamma_{22}$	$2\gamma_{23}$	$9\gamma_{24}$	
	Sep										1	$2\gamma_{16}$	γ_{17}	γ_{18}	γ_{19}	γ_{20}	$2\gamma_{21}$				$2\gamma_{22}$	$2\gamma_{23}$	$3\gamma_{24}$	
	O-D					4	4	4			1	γ_{16}												

Trial 5

For Trial 5, stock assignment for Hypotheses A and B are based on “stock70” assignment by STRUCTURE in *Data_NPM_190226_v3.csv*. The number of samples assigned to stock by sub-area is as follows. Table 7a of specifications details the assigned numbers by stock, sub-area, period and sex used to condition the trials.

Males	10E	11	1E	2C	6E	7CN	7CS	7E	7WR	8	9
J-stock	9	30	30	114	482	171	151	0	2	2	5
O-stock	1	35	1	27	3	625	308	44	76	223	470
Unassigned	1	0	1	2	10	22	18	0	2	4	12
Females											
J-stock	6	30	43	200	495	118	161	0	1	0	0
O-stock	0	33	0	27	5	273	314	5	8	20	52
Unassigned	1	2	1	2	7	7	11	0	0	3	2

Pink highlight: animals of a stock have not been sampled in a sub-area, but are allowed in that sub-area in the mixing matrices.

Green highlight: indicates sub-areas that differ in presence/absence in Trial 5 from the baseline trials.

- The distribution of J-stock whales is assumed to extend further in Trial 5 compared to the baseline, and are thus assumed to be found in sub-areas 7WR, 8 and 9 and by default therefore also in sub-area 7E.

Grey highlight: stock has been assigned to a sub-area, but is not modelled in that sub-area in the mixing matrices.

- No further extension in the distribution of O-stock whales from that assumed in the baseline is assumed in Trial 5 (e.g. into sub-areas 1E or 6E) due to the small assignments of O-stock whales for sub-areas 1E and 6E compared to the number of J-stock whales assigned to these sub-areas.

For Trial 5, stock assignment for Hypothesis E is based on “geneland.stock4” assignment by GENELAND in *Data_NPM_190226_v3.csv*. The number of samples assigned to stock by sub-area is as follows. Table 7a of specifications details the assigned numbers by stock, sub-area, period and sex used to condition the trials.

Males	10E	11	1E	2C	6E	7CN	7CS	7E	7WR	8	9
J-stock	8	14	31	96	492	21	4	0	0	0	0
P-stock	0	40	0	11	0	390	240	0	0	0	0
O-stock	0	1	0	0	0	308	91	42	77	217	478
Unassigned	0	8	0	20	0	55	111	0	0	0	0
Females											
J-stock	7	18	44	164	501	20	2	0	0	0	0
P-stock	0	24	0	11	0	219	312	0	0	0	0
O-stock	0	4	0	1	0	62	20	5	9	23	52
Unassigned	0	18	0	26	0	77	124	0	0	0	0

Appendix 3

Calculation of stock mixing proportions, including correction for “missing alleles”:

Unpooled results for sub-area 6W

C.L. de Moor

This appendix is based on de Moor (2011, 2014) and details the calculation of the stock mixing proportions by month and sex used for conditioning the 2013 *Implementation Simulation Trials* of western North Pacific common minke whales (Allison et al, 2014).

In calculating the mixing proportions in sub-area 6W, samples representative of ‘pure’ Y-stock and J-stock animals were taken as follows:

Stock	Location / months to define pure sample	Haplotypes Sample Size	Loci Sample Size
Y-stock	5 (all months)	58	58 58 58 58 58 56 58 58 54
J-stock	6E (all months)	392	392 392 392 392 392 392 392 392 392 (392 391 392 392 392)

Mixing proportions in sub-area 6W were calculated from 415 samples from bycatch data only.

Hyp B and E: Proportion of J mixing with Y		Sample Size	Proportion Haplotypes	SE	Sample Size (x11)	Proportion Loci	SE	
Jan-Mar	Males	83	0.555	0.142	83 with 81 in 11 th	0.745	0.050	
		37	0.449	0.253	37 with 36 in 1 st	0.963	0.083	
		41	0.749	0.243	41 with 40 in 8 th	0.926	0.062	
		43	0.534	0.245	43	0.787	0.080	
		21	0.830	0.38	21	0.788	0.089	
		16	1.000	0.004	16 with 15 in 11 th	0.726	0.137	
		20	0.533	0.335	20 with 18 in 11 th	0.475	0.107	
		97	0.629	0.140	97 with 96 in 7 th and 94 in 11 th	0.859	0.049	
Jan-Mar	Females	13	0.730	0.314	13 with 12 in 6 th	0.284	0.128	
		3	0.002	0.139	3	0.751	0.301	
		7	0.000	0.006	7	0.529	0.148	
		10	0.364	0.309	10	0.583	0.167	
		1	1.000	0.009	1	0.999	0.000	
		4	1.000	0.024	4	0.457	0.323	
		6	0.415	0.636	6 with 5 in 9 th	0.773	0.143	
		13	0.409	0.455	13 with 12 in 11 th	0.806	0.130	
Summary: all data		415	0.625	0.069	415 with 414 in 1 st , 6- 9 th and 406 in 11 th	0.776	0.109	
Pooled Data								
Jan-Mar	M F	96	0.584	0.131	96 with 95 in 6 th , 94 in 11 th	0.672	0.047	
Apr-Jun	M F	141	0.496	0.126	141 with 140 in 1 st , 8 th	0.812	0.04	
Jul-Aug	M F	42	1.000	0.004	42 with 41 in 11 th	0.749	0.077	
Sep-Dec	M F	136	0.593	0.123	136 with 135 in 7 th , 9 th , 130 in 11 th	0.761	0.04	

Notation:

In most cases samples are obtained from 16 loci. In sub-area 6W samples from the first 11 loci only were available to be used in the calculation of the mixing proportions, denoted by (x11) in the above table. In some cases there was a missing value in a sample at a particular loci. Thus, for example if the total sample size were 50, for one of the loci (the 10th) the sample size is 49. This is noted by saying e.g. “50 with 49 in 10th”.

References

- Allison, C., de Moor, C.L. and Punt, A.E. 2014. Report of the Scientific Committee. Annex D1. Report of the Working Group on the *Implementation Review* for Western North Pacific Common Minke Whales. Appendix 2. North Pacific minke whale *Implementation Simulation Trial* specifications. *J. Cetacean Res. Manage. (Suppl.)* 15:133-80.
- de Moor, C.L. 2011. Calculation of stock mixing proportions, including correction for ‘missing alleles’: unpooled results. Paper SC/D11/NPM4rev presented to the First Intersessional Workshop for the *Implementation Review* of western North Pacific common minke whales, 12-16 December 2011, Tokyo, Japan (unpublished). [Paper available from the Office of this Journal].
- de Moor, C.L. 2014. Report of the Scientific Committee. Annex D1. Report of the Working Group on the *Implementation Review* for Western North Pacific Common Minke Whales. Appendix 2. North Pacific minke whale *Implementation Simulation Trial* specifications. Adjunct 3. Calculation of stock mixing proportions, including correction for ‘missing alleles’: unpooled results. *J. Cetacean Res. Manage. (Suppl.)* 15:167-80.

Appendix 4

Method to derive the adjusted coefficient of variation for zero survey estimates

AE Punt

Simple case - the data are the number of observed whales and the sampling process is Poisson (for the case of one area):

$$LnL = \sum_y (n_y^{obs} \ln(\beta_y P_y) - \beta_y P_y) \quad (1)$$

where n_y^{obs} is the observed number of animals during the survey in year y , P_y is the true population size in year y , and β_y is the proportion of the area occupied that was sampled. For $\beta_y = 1$ this collapses to the standard Poisson likelihood.

Now consider the situation in which there is over-dispersion (e.g. clumping), one can account for this by defining an over-dispersed distribution for the data, i.e.

$$LnL > \sum_y (n_y^{obs} \ln(\beta_y P_y) - \beta_y P_y) / \alpha \quad (2)$$

where α is a measure of overdispersion (and would be greater than 1 for over-dispersed sampling). Justin provided the following formula for α :

$$\alpha = \sum_{y'} CV(P_{y'}^{obs}) n_{y'}^{obs} / \sum_{y'} 1 \quad (3)$$

where the summation is over years for which there is a CV for the abundance estimate and a value for the number of sightings.

To derive 3, one estimator for α is:

$$\alpha = \sum_y \frac{\text{var(observed}_y\text{)}}{\text{var(expected}_y\text{)}} / \sum_y 1 \quad (4)$$

where $\text{var(observed}_y\text{)} = CV^2(P_y)(n_y^{obs})^2$ and $\text{var(expected}_y\text{)} = n_y^{obs}$ (under the Poisson assumption) so that

$$\frac{\text{var(observed}_y\text{)}}{\text{var(expected}_y\text{)}} \sim \frac{(n_y^{obs})^2 CV^2(n_y^{obs})}{E(n_y^{obs})} \cong \frac{(n_y^{obs})^2 CV^2(P_y)}{n_y^{obs}} = n_y^{obs} CV^2(P_y) \quad (5)$$

which is close to, but not identical to, 3. An alternative estimate for α would be:

$$\alpha = \frac{\sum_y \text{var(observed}_y\text{)}}{\sum_y \text{var(expected}_y\text{)}} \cong \frac{\sum_y (n_y^{obs})^2 CV^2(P_y)}{\sum_y n_y^{obs}} \quad (6)$$

Equation 6 would (I suspect) be more robust to odd outlying estimates of CV.